arXiv:1111.3392v1 [math.AG] 14 Nov 2011

NONCOMMUTATIVE MIRROR SYMMETRY FOR PUNCTURED SURFACES

RAF BOCKLANDT

ABSTRACT. In [7] Abouzaid, Auroux, Efimov, Katzarkov and Orlov showed that the wrapped Fukaya Categories of punctured spheres are derived equivalent to the categories of singularities of a superpotential on certain crepant resolutions of toric 3 dimensional singularities. We generalize this result to other punctured Riemann surfaces and reformulate it in terms of certain noncommutative algebras coming from dimer models. In particular, given any consistent dimer model we can look at a subcategory of noncommutative matrix factorizations and show that this category is $\mathtt{A}_\infty$-isomorphic to a subcategory of the wrapped Fukaya category of a punctured Riemann surface. The connection between the dimer model and the punctured Riemann surface then has a nice interpretation in terms of a duality on dimer models. Finally, we tie this result to the classical commutative mirror symmetry.

1. INTRODUCTION

Originally homological mirror symmetry was developed by Kontsevich [30] as a framework to explain the similarities between the symplectic geometry and algebraic geometry of certain Calabi-Yau manifolds. More precisely its main conjecture states that for any compact Calabi-Yau manifold with a complex structure X, one can find a mirror Calabi-Yau manifold X' equiped with a symplectic structure such that the derived category of coherent sheaves over X is equivalent to the zeroth homology of the Fukaya category of X'. The latter is a category that represents the intersection theory of Lagrangian submanifolds of X'.

$$\mathtt{D}^b\mathtt{Coh}\, X \sim \mathtt{H}^0\mathtt{Fuk}X'$$

Over the years it turned out that this conjecture is part of a set of equivalences which are much broader than the compact Calabi-Yau setting [26, 20, 1, 5, 6]. Removing the compactness or Calabi-Yau condition often makes the mirror a singular object, which physicists call a Landau-Ginzburg model [36, 37].

A Landau-Ginzburg model (X, W) is a pair of a smooth space X and a complex valued function $W : X \to \mathbb{C}$, which is called the potential. On the algebraic side we associate to it the dg-category of matrix factorizations $\mathtt{MF}(X, W)$. Its objects are diagrams $P_0 \rightleftarrows P_1$ where P_i are vector bundles and the composition of the maps results in multiplication with W. The morphisms are morphisms between these vector bundles equiped with a natural differential.

On the symplectic side we demand that the Lagrangians have their boundary fixed on $W^{-1}(0)$. On the other hand if X' is noncompact we need to tweak the notion of the Fukaya category, by imposing certain conditions on the behaviour of the Langrangians near infinity and using a Hamiltonian flow to adjust the intersection theory. This gives us the notion of a wrapped Fukaya category [2].

In [7] Abouzaid, Auroux, Efimov, Katzarkov and Orlov proved an instance of mirror symmetry between such objects. On the symplectic side they considered a sphere with k punctures and on the algebraic side they considered a special Landau Ginzburg model on a certain toric quasiprojective noncompact Calabi Yau threefold and they proved an equivalence between the derived wrapped Fukaya category of the former and the derived category of matrix factorizations of the latter.

In this paper we aim to generalize their result to all Riemann surfaces with $k \geq 3$ punctures. On the algebraic side though we will not construct classical Landau-Ginzburg models but instead we will look at noncommutative Landau-Ginzburg models [38]. This means that we replace the commutative space X with a noncommutative Calabi-Yau algebra A and the potential will be a central element. The category of matrix factorizations needs also an adjustment: instead of vector bundles we must take projective modules.

The Calabi-Yau algebras under consideration will come from dimer models, which are certain quivers embedded in a Riemann surface. Such algebras, known as Jacobi algebras, also have a canonical choice of potential ℓ coming from the faces in which the quiver splits the Riemann surface.

The result we obtain is that for any consistent dimer model Q, we can find a full subcategory $\mathtt{mf}(\mathrm{Q})$ of the category of all matrix factorizations of its Jacobi algebra $\mathtt{MF}(\mathtt{Jac}\,\mathrm{Q}, \ell)$ which is $\mathtt{A}_\infty$-isomorphic to a full subcategory $\mathtt{fuk}(\reflectbox{Q})$ of the wrapped Fukaya category $\mathtt{Fuk}(S)$ of a punctured Riemann surface S. The category $\mathtt{fuk}(\reflectbox{Q})$ is constructed using a new dimer model $\reflectbox{Q}$, embedded in the closure of S such that its vertices are the punctures. The new dimer can be obtained explicitly from the original dimer by a process called dimer duality.

We illustrate how our viewpoint and the approach in [7] fit together. The simplest example gives us an equivalence between the sphere with 3 punctures $SS^{\bullet 3}$ and the Landau-Ginzburg model $(\mathbb{C}^3, xyz)$. Commutatively this corresponds to the category of singularities of the three standard planes in affine 3-space. Noncommutatively[1] we can see this as a dimer model on the torus with a superpotential ℓ coming from the faces.


Commutative version
$\mathtt{H}^0\mathtt{MF}(\mathbb{C}^3, xyz)$
$\sim$
$\mathtt{H}^0\mathtt{Fuk}(SS^{\bullet 3})$
Noncommuative version
$\mathtt{mf}(\mathrm{Q})$
$\cong_\infty$
$\mathtt{fuk}(\reflectbox{Q})$
y
x
z
y
x
z
1
x
1
y
z
y
1
x
1
1
y
3
y
z
x
2
x
1

On the right hand side Q is embedded in a torus while its dual, $\reflectbox{Q}$, sits in a sphere an its 3 vertices are the 3 punctures in the commutative picture. The dual can be obtained by flipping over the clockwise faces, reversing their arrows and gluing everything back again.

By construction $\mathtt{fuk}(\reflectbox{Q})$ will generate the full wrapped Fukaya category. On the algebra side it is not completely clear whether $\mathtt{mf}(\mathrm{Q})$ generates the full category of matrix factorizations $\mathtt{MF}(\mathtt{Jac}\,\mathrm{Q}, \ell)$ because unless Q sits inside a torus, the Jacobi algebra is not Noetherian. However if Q does sit inside a torus, we will show that $\mathtt{mf}(\mathrm{Q})$ indeed generates $\mathtt{MF}(\mathtt{Jac}\,\mathrm{Q}, \ell)$. A result of Ishii and Ueda [23] also implies that $\mathtt{MF}(\mathtt{Jac}\,\mathrm{Q}, \ell) \cong \mathtt{MF}(\tilde{X}, \ell)$ where $\tilde{X}$ is a crepant resolution of the center $\mathsf{Spec} Z(\mathtt{Jac}\,\mathrm{Q})$, so we recover the commutative result.

The outline of the paper is as follows: first we review some of the basics of $\mathtt{A}_\infty$-structures, quivers, dimer models and some algebras associated to them. We combine these subjects to look at $\mathtt{A}_\infty$-structures on certain dimer models, called rectified dimers. We classify some of these $\mathtt{A}_\infty$-structures and then we turn to both sides of mirror symmetry. First we show that the Fukaya category associated to any polyhedral subdivision of a Riemann-surface gives rise to one of the $\mathtt{A}_\infty$-structures we considered. Secondly we show that matrix factorizations of a consistent dimer model also give rise to such an $\mathtt{A}_\infty$-structure and finally we tie the two sides together by constructing an explicit duality on dimer models. We end with a discussion about the connection between the commutative results in [7] and the noncommutative geometry we employed. In view of readability of the paper, we deferred the proof of the classification of $\mathtt{A}_\infty$-structures to an appendix.

---

[1] Note that in this example the Jacobi algebra $\mathtt{Jac}\,\mathrm{Q} \cong \mathbb{C}[X, Y, Z|$ is not noncommutative, but for all other dimer models it is.

2. $\mathtt{A}_\infty$-CATEGORIES

In this section we will introduce the basics of $\mathtt{A}_\infty$-categories. For more information we refer to [27, 32].

An *$\mathtt{A}_\infty$-category $\mathtt{C}$ with degrees in $\mathbb{Z}_2$* consists of the following data[2]

- a set of objects $\mathtt{Ob}\,\mathtt{C}$,
- for each pair of object $X, Y \in \mathtt{Ob}\,\mathtt{C}$ a complex $\mathbb{Z}_2$-graded vector space $\mathtt{Hom}_\mathtt{C}(X, Y)$,
- for each sequence of $n+1$ objects $X_0, \dots X_k \in \mathtt{Ob}\,\mathtt{C}$ a multilinear map μ_k of degree $2-k$

  $$\mu_k : \mathtt{Hom}_\mathtt{C}(X_1, X_0) \otimes \cdots \otimes \mathtt{Hom}_\mathtt{C}(X_k, X_{k-1}) \to \mathtt{Hom}_\mathtt{C}(X_k, X_0)$$

  (if there is no confusion we drop the subscript and just write μ)

such that the identities

$$[\mathrm{M_k}] \quad \sum_{s+l+t=k} (-1)^\sigma \mu(f_1, \dots, f_s, \mu(f_{s+1}, \dots, f_{s+l}), f_{s+l+1}, \dots, f_k) = 0$$

hold for any sequence of homogeneous homomorphisms $X_0 \xleftarrow{f_1} X_1 \cdots X_{k-1} \xleftarrow{f_k} X_k$. In this expression the sign takes the form $\sigma = s + lt + (2-l)(\deg f_1 + \cdots + \deg f_k)$. For $k=1$, the identity becomes $\mu\mu(f) = 0$, so each hom-space can be considered as a complex with $d : f \mapsto \mu(f)$.

An $\mathtt{A}_\infty$-category is called *strictly unital* if for every object X there is an element $\mathbf{1}_X \in \mathtt{Hom}_\mathtt{C}(X, X)$ of degree 0 such that

- $\mu(a, \mathbf{1}_X) = a$ if a is a homomorphism with source X and $\mu_2(1_X, a) = a$ is a homomorphism with target X.
- $\mu(a_1, \dots, a_n) = 0$ if $n \neq 2$ and one of the $a_i = \mathbf{1}_X$.

If $\mu_i = 0$ for all $i \neq 2$ a strictly unital $\mathtt{A}_\infty$-category is the same as a $\mathbb{Z}_2$-graded ordinary category. If $\mu_i = 0$ for all $i \geq 3$ we get a dg-category.

Aside 2.1. Just like for ordinary categories we define an $\mathtt{A}_\infty$-algebra as an $\mathtt{A}_\infty$-category with one object and identifying the algebra with the hom-space from this object to itself. We can turn any $\mathtt{A}_\infty$-category into an $\mathtt{A}_\infty$-algebra by taking the direct sum of all hom spaces in the original category and extending the products multilinearly and setting products of maps that do not concatenate zero.

An *$\mathtt{A}_\infty$-functor* $\mathcal{F} : \mathtt{A} \to \mathtt{B}$ between two $\mathtt{A}_\infty$-categories consists of

- a map $\mathcal{F}_0 : \mathtt{Ob}\,\mathtt{A} \to \mathtt{Ob}\,\mathtt{B}$
- for each sequence of $k+1$ objects $X_0, \dots X_k \in \mathtt{Ob}\,\mathtt{A}$ a linear map of degree $1-k$

  $$\mathcal{F}_k : \mathtt{Hom}_\mathtt{C}(X_1, X_0) \otimes \cdots \otimes \mathtt{Hom}_\mathtt{C}(X_k, X_{k-1}) \to \mathtt{Hom}_\mathtt{C}(\mathcal{F}_0 X_k, \mathcal{F}_0 X_0)$$

  (if there is no confusion we drop the subscript and just write $\mathcal{F}$)

subject to the following identities

$$[\mathrm{F_k}] \begin{cases} \sum_{0 \leq s < t \leq k} \pm \mathcal{F}(f_1, \dots, f_s, \mu(f_{s+1}, \dots, f_t), f_{t+1}, \dots, f_k) \\ = \sum_r \sum_{1 \leq i_1 \leq \cdots \leq i_r < k} \pm \mu(\mathcal{F}(f_1, \dots, f_{i_1}), \dots, \mathcal{F}(f_{i_{r+1}}, \dots, f_k)) \end{cases}$$

for any sequence of homogeneous homomorphisms $X_0 \xleftarrow{f_1} X_1 \cdots X_{k-1} \xleftarrow{f_k} X_k$. An $\mathtt{A}_\infty$-functor is called *strict* if $\mathcal{F}_i = 0$ for $i > 1$, it is called an *isomorphism* if $\mathcal{F}_0$ is a bijection and all $\mathcal{F}_1$ are isomorphisms and it is called a *quasi-isomorphism* if $\mathcal{F}_0$ is a bijection and the $\mathcal{F}_1$ induce isomorphisms on the level of the homology of $d = \mu_1$.

---

[2] Everything in this section can be generalized to $\mathtt{A}_\infty$-categories with degrees in $\mathbb{Z}$ or any other cyclic group.

2.1. **Minimal models.** An A_∞-category is called *minimal* if $\mu_1 = 0$. Note that unlike general $\mathtt{A}_\infty$-categories, minimal $\mathtt{A}_\infty$-categories are also genuine categories if we put $f_1 f_2 := \mu(f_1, f_2)$ and forget the higher μ's. This is because in this case $[\mathrm{M}_3]$ becomes the standard associativity identity.

An $\mathtt{A}_\infty$*-structure on an ordinary* $\mathbb{Z}_2$*-graded* $\mathbb{C}$*-linear category* $\mathtt{C}$ is a set of multiplications μ that turn $\mathtt{C}$ into a minimal $\mathtt{A}_\infty$-category such that as an ordinary category it is identical to the category structure of $\mathtt{C}$.

Theorem 2.2 (Kadeishvili)**.** [25] *Let* $\mathtt{C}$ *be an* A_∞*-category and denote by* $\mathtt{H}\,\mathtt{C}$ *the 'category' with the same objects but as hom-spaces the homology of the hom-spaces in* $\mathtt{C}$*. There is an* $\mathtt{A}_\infty$*-structure on* $\mathtt{H}\,\mathtt{C}$ *such that there is an object-preserving quasi-isomorphism between* $\mathtt{H}\,\mathtt{C}$ *and* $\mathtt{C}$*. This* $\mathtt{A}_\infty$*-structure on* $\mathtt{H}\,\mathtt{C}$ *is unique up to object-preserving* $\mathtt{A}_\infty$*-isomorphism.*

How do we construct this new $\mathtt{A}_\infty$-structure? In order to do this we use a graphical method from [31]. Set $d = \mu_1$ and for each $\mathtt{Hom}_{\mathtt{C}}(X, Y)$ choose a map h of degree -1 on A such that

$$h^2 = 0 \text{ and } dhd = d.$$

We will call this map a codifferential. The map $\pi := 1 - dh - hd$ is a projection and we can identify $\mathrm{Im}\pi$ with $\mathtt{Hom}_{\mathtt{H}\,\mathtt{C}}(X, Y)$ because $d\pi = \pi d = 0$ and if $dx = 0$ then $\pi x = x - d(hdx)$. Let ι be the embedding $\mathtt{Hom}_{\mathtt{H}\,\mathtt{C}}(X, Y) = \mathrm{Im}\pi \subset \mathtt{Hom}_{\mathtt{C}}(X, Y)$.

Given a rooted tree $\mathcal{T}$ with $k + 1$ leaves we can define a multilinear map

$$m^{\mathcal{T}} : \mathtt{H}\,\mathtt{Hom}_{\mathtt{C}}(X_1, X_0) \otimes \cdots \otimes \mathtt{H}\,\mathtt{Hom}_{\mathtt{C}}(X_k, X_{k-1}) \to \mathtt{H}\,\mathtt{Hom}_{\mathtt{C}}(\mathcal{F}_0 X_k, \mathcal{F}_0 X_0)$$

by interpreting every leaf as the map ι, every internal node as a map $h\mu$ and the root as $\pi\mu$.



$$\Longrightarrow\ m^{\mathcal{T}}(f_1, \ldots, f_5) = \pi\mu(h\mu(f_1, f_2), h\mu(f_3, f_4, f_5)).$$

The new multiplication $\mu_n^{\mathtt{H}}$ is then defined as the sum $\sum_{\mathcal{T}} m^{\mathcal{T}}$ over all rooted with n leaves for $n > 1$ and $\mu_1^{\mathtt{H}} = 0$.

2.2. **Twisted completion.** Given an $\mathtt{A}_\infty$-category $\mathtt{C}$ we define a *twisted object* [27] as a pair (M, δ), where $M \in \mathbb{N}[\mathtt{Ob}\,\mathtt{C} \times \mathbb{Z}_2]$ is a formal sum of objects shifted by elements in $\mathbb{Z}_2$. We will write such a sum as $v_1[i_1] \oplus \cdots \oplus v_k[i_k]$ where the v_j are objects and the i_j shifts. The map δ is an uppertriangular $k \times k$-matrix with entries $\delta_{st} \in \mathtt{Hom}_{\mathtt{C}}(v_{i_t}, v_{i_s})$ of degree 1 and subject to the identity

$$\sum_{n=1}^{\infty} (-1)^{\frac{n(n-1)}{2}} \mu_n(\delta, \ldots, \delta) = 0$$

where we extended μ_n to matrices in the standard way.

The homomorphism space between two such objects (M, δ) and (M', δ') is given by

$$\bigoplus_{r,s} \mathtt{Hom}(v_r, v'_s)[i_r - i_s]$$

which we equip with an $\mathtt{A}_\infty$-structure as follows:

$$\mu(f_1, \ldots, f_n) := \sum_{t=0}^{\infty} \sum_{i_0 + \cdots + i_n = t} \pm\mu(\underbrace{\delta, \ldots, \delta}_{i_0}, f_1, \underbrace{\delta, \ldots, \delta}_{i_1}, \ldots, f_n, \underbrace{\delta, \ldots, \delta}_{i_n}).$$

The $\pm$-sign is calculated by multiplying with a factor $(-1)^{n+t-k}$ for each δ in the expression on position k.

The $\mathtt{A}_\infty$-category of twisted objects and their homomorphism spaces is denoted by $\mathtt{Tw}\,\mathtt{C}$. It also has a minimal model, which we denote by $\mathtt{H}\,\mathtt{Tw}\,\mathtt{C}$. Note that because it is a minimal model, $\mathtt{H}\,\mathtt{Tw}\,\mathtt{C}$ is a genuine category.

Aside 2.3. If $\mathtt{A}$ is a genuine $\mathbb{C}$-linear category with a finite number of objects, such that the $\mathtt{Hom}_A(X,Y)$ are finite dimensional and contain no nontrivial idempotents, then we can consider $\mathtt{A}$ as a path algebra of a quiver with relations. We can construct the derived category $\mathtt{DModA}$ and look at the smallest triangulated subcategory generated by $\mathtt{A}$ as a module over itself. We can construct a $\mathbb{Z}$-graded category from this by putting $\mathtt{Hom}(X,Y) := \oplus_{i\in\mathbb{Z}}\mathtt{Hom}_{\mathtt{DModA}}(X,Y[i])$.

On the other hand we can view $\mathtt{A}$ as an $\mathtt{A}_\infty$-category with degrees in $\mathbb{Z}$, by putting all of $\mathtt{Hom}_{\mathtt{A}}(X,Y)$ in degree 0. It makes sense to look at $\mathtt{H}\,\mathtt{Tw}\,\mathtt{A}$ and it turns out that this category is equivalent to the category we defined above (see [27]). In this light $\mathtt{H}\,\mathtt{Tw}\,\mathtt{C}$ can be seen as a useful generalization of the derived category of an algebra.

3. EMBEDDED QUIVERS AND DIMERS

As usual a *quiver* Q is a finite (or locally finite) oriented graph. We denote the set of vertices by Q_0, the set of arrows by Q_1 and the maps h, t assign to each arrow its head and tail. A *nontrivial path* p is a sequence of arrows $a_k \cdots a_0$ such that $t(a_i) = h(a_{i+1})$. We write $p[i]$ to denote the arrow a_i and we set $h(p) = h(p[k])$ and $t(p) = t(p[0])$.

A *trivial path* is just a vertex. A path p is called *cyclic* if $h(p) = t(p)$ and the equivalence class of a cyclic path under cyclic permutation is called a *cycle*.

The *path category* $\mathbb{C}Q$ is the category with as objects the vertices, as homomorphisms linear combinations of paths and as composition concatenation.

Given a quiver Q, one can construct its double, $\bar{Q}$, which has the same number of vertices but for every arrow $a \in Q_1$ we add an extra arrow a^{-1} with $h(a) = t(a^{-1})$ and $t(a) = h(a^{-1})$. The *weak path category* of Q is the following quotient

$$\mathbb{C}\hat{Q} := \frac{\mathbb{C}Q}{\langle aa^{-1} = h(a), a^{-1}a = t(a) | a \in Q_0\rangle}$$

A *weak path in* Q is a path in $\bar{Q}$, viewed as a homomorphism in the weak path category. We also speak of weak arrows and cycles and if we want to stress that a path or cycle is not weak we will call it real.

A quiver is called *embedded in a compact orientable surface without boundary* S, if Q_0 is a discrete subset of S and every arrow is a smooth embedding $a : [0,1] \to S$ such that $h(a) = a(1)$ and $t(a) = a(0)$ and different arrows only intersect in end points. We identify a^{-1} with the map a in reverse direction. We also denote the surface in which the quiver embeds by $|Q|$.

If Q is a quiver embedded in S and $\pi : \tilde{S} \to S$ is the universal cover of S, we can construct a new quiver $\tilde{Q}$ which is called the *universal cover of* $\tilde{Q}$. Its vertices are all points in $\tilde{S}$ that map onto vertices in Q, and its arrows are maps that push forward to arrows in Q. Note that given a vertex $v \in Q_0$ and a lift in $\tilde{v} \in \tilde{Q}_0$, we can lift every path in Q starting in v to a unique path starting in $\tilde{v}$.

We say that the quiver Q *splits the surface* S if the complement of the quiver consists of a disjoint union of open disks. Each of these disks is bounded by a weak path c that goes around it in counterclockwise direction. Such a cycle is called a boundary cycle and we collect these cycles in a set $\hat{Q}_2^+$.

A *dimer model*[17, 14, 16] is a quiver that splits a surface for wich every boundary cycle is either real or the inverse of a real cycle. The real cycles that are boundary cycles are called the positive cycles and are grouped in the set $Q_2^+ := \hat{Q}_2^+ \cap \mathbb{C}Q$. Negative cycles are those for which the inverse is a boundary cycle and we set $Q_2^- := (\hat{Q}_2^+)^{-1} \cap \mathbb{C}Q$. Note that the orientability of the surface implies that every arrow will be contained in one positive cycle and one negative.

Example 3.1. We give 4 examples of embedded quivers that split a surface. The first 3 are embedded in a torus, the last in a double torus. Arrows and vertices with the same label are identified.



The last 3 are dimer models.

The *Jacobi category of a dimer model* is the quotient of the path category by the ideal generated by relations of the form $r_a := r_+ - r_-$ where $r_+ a \in \mathrm{Q}_2^+$ and $r_- a \in \mathrm{Q}_2^-$ for some arrow $a \in \mathrm{Q}_1$:

$$\mathsf{Jac}\,(\mathrm{Q}) := \frac{\mathbb{C}\mathrm{Q}}{\langle r_a | a \in \mathrm{Q}_1 \rangle}.$$

For the last dimer model in example 3.1 we have $\mathrm{Q}_2^+ = \{abxcd\}$ and $\mathrm{Q}_2^- = \{baxdc\}$ and

$$\mathsf{Jac}\,(\mathrm{Q}) := \frac{\mathbb{C}\langle a,b,c,d,x \rangle}{\langle bxcd - xcdb, xcda - axdc, cdab - dcba, dabx - baxd, abxc - cbax \rangle}.$$

Aside 3.2. The reason these Jacobi algebras are important is because they appear as noncommutative analogues of Calabi-Yau manifolds. A compact 3-Calabi-Yau manifold can be defined as a smooth variety X which has the following duality

$$\mathsf{Ext}^i_X(\mathscr{F},\mathscr{G}) = \mathsf{Ext}^{3-i}_X(\mathscr{G},\mathscr{F})^*$$

for all coherent sheafs $\mathscr{F},\mathscr{G}$ on X.

Similarly a 3-Calabi-Yau algebra [15] can be defined such that it has a similar duality

$$\mathsf{Ext}^i_X(M,N) = \mathsf{Ext}^{3-i}_X(N,M)^*$$

for all finite dimensional left A-modules.

Not every Jacobi algebra will be Calabi-Yau, this will depend on the structure of the dimer model. To characterize such dimer models we introduce a notion of consistency. Several different notions are available in the literature [21, 11, 12, 35, 24, 10] but we will restrict to one: zigzag consistency.

Fix a dimer model Q and its universal cover $\tilde{\mathrm{Q}}$. For any arrow $\tilde{a} \in \tilde{\mathrm{Q}}_1$ we can construct its *zig ray* $\mathcal{Z}^+_{\tilde{a}}$. This is an infinite path

$$\dots \tilde{a}_2 \tilde{a}_1 \tilde{a}_0$$

such that $\tilde{a}_0 = \tilde{a}$ and $\tilde{a}_{i+1}\tilde{a}_i$ sits in a positive cycle if i is even and in a negative cycle if i is odd. Similarly the *zag ray* $\mathcal{Z}^-_{\tilde{a}}$ is the path where $\tilde{a}_{i+1}\tilde{a}_i$ sits in a positive cycle if i is odd and in a negative cycle if i is even. The projection of a zig or a zag ray down to Q will give us a cyclic path because Q is finite. Such a cyclic path will be called a *zigzag cycle*. A dimer model is called *zigzag consistent* if for every arrow $\tilde{a}$ the zig and the zag ray only meet in $\tilde{a}$:

$$\mathcal{Z}^-_{\tilde{a}}[i] = \mathcal{Z}^+_{\tilde{a}}[j] \implies i = j = 0.$$

In example 3.1 the second and fourth quiver are consistent, while the third quiver is not because $\mathcal{Z}^-_{\tilde{x}}[3] = \mathcal{Z}^+_{\tilde{x}}[3] = \tilde{y}$. Note that a dimer model on a sphere is never zigzag consistent as its universal cover is finite.

Theorem 3.3. [10] *If a dimer model is zigzag-consistent then its Jacobi Algebra is 3-Calabi-Yau.*

4. RECTIFIED DIMERS AND GENTLE CATEGORIES

Given a quiver Q that splits the surface we construct a dimer model, which we call *the rectified dimer* RQ.

- The vertices of RQ are the centres of the arrows in Q
$$\mathrm{RQ}_0 := \{v_a := a(1/2) | a \in Q_1\}.$$
- The arrows of RQ are segments $\alpha = \overleftarrow{v_a v_b}$ connecting the centres of two (weak) arrows that follow each other clockwise in one of the boundary cycles.

It is easy to check that this is indeed a dimer model. Each of the boundary cycles of Q will give a positive cycle in RQ and each of the vertices in RQ will give us a negative cycle. In Euclidean geometry the process of cutting of the vertices of a polyhedron at the midpoints of the edges is called rectification, hence the name.

To each of the arrows we assign a $\mathbb{Z}_2$-degree. If abr is a boundary cycle of Q and a and b are both real or both weak we give the arrow $\alpha = \overleftarrow{v_a v_b}$ degree 1. Otherwise we give it degree 0. Note that if Q is already a dimer then all arrows in RQ have degree 1.

Example 4.1. Below are two examples of rectification. In the first we rectify a quiver on a torus with one vertex and two loops to obtain a dimer model with two vertices and four arrows. In the second example we rectify a tetrahedron to obtain an octahedron.





To indicate the grading we marked the degree zero arrows. The unmarked arrows all have degree 1.

Instead of associating to this dimer its Jacobi category, we will look at a different category. The *gentle category of a rectified dimer* is the quotient of the path category by the ideal generated by relations of the form $\alpha\beta$ where α and β are consecutive arrows of a positive cycle.

$$\mathtt{Gtl}\,(\mathrm{RQ}) := \frac{\mathbb{C}\mathrm{RQ}}{\langle \alpha\beta | \exists \rho : \alpha\beta\rho \in \mathrm{RQ}_2^+ \rangle}$$

Remark 4.2. The gentle category of a rectified dimer is a generalization of the gentle algebra of a triangulation which was introduced by Assem, Brüstle, Plamondon and Charbonneau-Jodoin in [3]. If the original quiver that splits the surface comes from a triangulation, these two algebras coincide. One can easily check that $\mathtt{Gtl}\,(\mathrm{RQ})$ is a gentle algebra in the sense of [3] and [4].

We end this section with a lemma on the structure of $\mathtt{Gtl}\,(\mathrm{RQ})$ that will be useful later on.

Lemma 4.3. *If v, w are two vertices in the rectified dimer RQ then*

$$v\mathtt{Gtl}\,(\mathrm{RQ})w = \begin{cases} p\mathbb{C}[c] & \text{if } c = vpwq \text{ is a negative cycle} \\ \mathbb{C}[c_1, c_2]/(c_1c_2) & v = w \text{ and } c_1, c_2 \text{ are the two negative cycles through } v \\ 0 & \text{otherwise} \end{cases}$$

Moreover all negative cycles have even degree.

Proof. Any nonzero nontrivial path can only be extended in one way to a nonzero path that is one arrow longer: one must add the arrow that sits in the same negative cycle as the last arrow. So, if p is a nonzero path then all its arrows must be contained in one negative cycle. This implies the statement above.

The degree of a negative cycle is even because if on changes the direction in one of the arrows in the orginal dimer, the degree of the cycle doesn't change (2 rectified arrows change degree). If all original arrows arrive in the original vertex representing the negative cycle then all rectified arrows in this cycle have zero degree. ☐

We will now describe a specific $\mathtt{A}_\infty$-structure on $\mathtt{Gtl}\,(\mathrm{RQ})$, which can be constructed inductively. For any sequence of paths $p_1, \dots, p_k$ and any cycle $b_1 \dots b_l \in \mathrm{R}Q_2^+$ with $h(b_1) = t(p_i)$ we set

$$\mu(p_1, \dots, p_i b_1, b_2, \dots, b_{l-1}, b_l p_{i+1}, \dots, p_k) := (-1)^s \mu(p_1, \dots, p_k).$$

with sign convention $s = l(p_1 + \dots + p_i + k - i)$.



For $k > 2$ we set $\mu(q_1, \dots, q_k) = 0$ if we cannot perform any reduction of the form above and for $k = 2$ we use the ordinary product on $\mathtt{Gtl}\,(\mathrm{R}Q)$. This $\mathtt{A}_\infty$-structure is an example of a broader class of $\mathtt{A}_\infty$-structures μ^κ, parametrized by maps $\mathrm{Q}_2^+ \times \mathbb{N} \to \mathbb{C}$, introduced in the appendix. In particular $\mu = \mu^\kappa$ with $\kappa(c, i) = 1$ if $i = 1$ and $\kappa(c, i) = 0$ if $i > 1$.

Theorem 4.4.

1. *μ defines an $\mathtt{A}_\infty$-structure on $\mathtt{Gtl}\,(\mathrm{RQ})$.*
2. *If μ is an $\mathtt{A}_\infty$-structure on $\mathtt{Gtl}\,(\mathrm{RQ})$ such that for every cycle $c = a_1 \dots a_l \in \mathrm{RQ}_2^+$ there is a $\lambda_c \in \mathbb{C}$ such that*

$$\mu(a_i, \dots, a_j) = \begin{cases} \lambda_c h(a_i) & j - i + 1 = 0 \mod l \\ 0 & j - i + 1 \neq 0 \mod l \end{cases}$$

then μ is $\mathtt{A}_\infty$-isomorphic to μ.

Proof. This is a combination of theorem 9.9, lemma 9.10 and theorem 9.12. Proofs of these results can be found in the appendix [**?**]. ☐

Different quivers in the same surface will give different gentle $\mathtt{A}_\infty$-categories, but these categories are closely related. In fact if we go to the twisted completion the difference disappears.

Lemma 4.5. *Suppose Q is an embedded quiver that splits the surface $|\mathrm{Q}|$ and a is one of the arrows of Q. If Q' is the new quiver obtained by changing the direction of a (i.e. swapping its head and tail) then $\mathtt{Tw}\,\mathtt{Gtl}\,(\mathrm{RQ}), \mu$ and $\mathtt{Tw}\,\mathtt{Gtl}\,(\mathrm{RQ}'), \mu$ are isomorphic $\mathtt{A}_\infty$-categories.*

Proof. Let v_0 be the object in $\mathtt{Gtl}\,(\mathrm{RQ})$ corresponding to the arrow a we want to reverse. We denote the corresponding object in $\mathtt{Gtl}\,(\mathrm{RQ}')$ as v_0'. All other objects we denote by v_i in both categories. We now define a strict functor $\mathcal{F} : \mathtt{Tw}\,\mathtt{Gtl}\,(\mathrm{R}Q) \to \mathtt{Tw}\,\mathtt{Gtl}\,(\mathrm{R}Q')$:

- $\mathcal{F}_0(v_0[0]^{\oplus m_0} + v_0[1]^{\oplus m_1} + \mathrm{rest}, \delta) = (v_0'[1]^{\oplus m_0} + v_0'[0]^{\oplus m_1} + \mathrm{rest}, \delta)$
- $\mathcal{F}_1(f) = f$

It is easy to check that the degrees match up and that this is an isomorphism. ☐

Lemma 4.6. *Suppose Q is an embedded quiver that splits the surface $|\mathrm{Q}|$. Suppose that $a_1 \dots a_k$ is a boundary cycle and let b be a new arrow in this face connecting $h(a_1)$ and $h(a_i)$ with $2 < i < k - 1$. Denote the quiver obtained by adding b to Q as Q', then $\mathtt{H}\,\mathtt{Tw}\,\mathtt{Gtl}\,(\mathrm{RQ})$ and $\mathtt{H}\,\mathtt{Tw}\,\mathtt{Gtl}\,(\mathrm{RQ}')$ are equivalent as A_∞-categories.*

Proof. Let v_0 be the object in $\mathtt{Gtl}\,(\mathrm{R}Q')$ corresponding to the arrow b we want to add. Denote the object corresponding to the arrow a_j by v_j in both categories. We use the Greek letters α_j to denote he arrow between v_{j+1} and v_j and the arrow β_0 connects v_1 with v_0 and β_i connects v_0 to v_i.

We can identify v_0 with the object

$$w = \left(v_1[1] + \cdots + v_i[1], \delta := \begin{pmatrix} 0 & \alpha_1 & & \\ & \ddots & \ddots & \\ & & & \alpha_{i-1} \\ & & & 0 \end{pmatrix} \right).$$

We have a map $f_1 : v_0 \to w$ given by $\begin{pmatrix} \beta_0 & 0 & \cdots & 0 \end{pmatrix}^\top$ and a map $f_2 : w \to v_0$ given by $\begin{pmatrix} 0 & \cdots & 0 & \beta_i \end{pmatrix}$.

These maps are each other's inverses in $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ}')$. Clearly $\mu_2(f_1, f_2) = \mu(f_1, \delta, \dots, \delta, f_2) = \mu(\beta_0, \alpha_1, \dots, \alpha_i, \beta_i) = \mathbf{1}_{v_0}$. On the other hand we have

$$\begin{aligned} \mu(f_2, f_1) &= \mu(f_2, f_1, \delta, \dots, \delta) + \mu(\delta, f_2, f_1, \delta, \dots, \delta) + \cdots + \mu(\delta, \dots, \delta, f_2, f_1) \\ &= \begin{pmatrix} \mu(\alpha_1,\dots,\alpha_{i-1},\beta_i,\beta_0) & & & \\ & \mu(\alpha_2,\dots,\alpha_{i-1},\beta_i,\beta_0,\alpha_1) & & \\ & & \ddots & \\ & & & \mu(\beta_i,\beta_0,\alpha_1,\dots,\alpha_{i-1}) \end{pmatrix} \\ &= \begin{pmatrix} v_1 & & & \\ & v_2 & & \\ & & \ddots & \\ & & & v_i \end{pmatrix} = \mathbf{1}_w \end{aligned}$$

Because the twisted completion is itself closed under twisted completion, $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ}')$ is a full subcategory of $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ})$. By construction the latter is also a full subcategory of the former, so they are equivalent. □

Corollary 4.7. *As an* $\mathtt{A}_\infty$*-category* $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ}), \mu$ *only depends on the genus of the surface and the number of the vertices of* Q.

Proof. By adding enough arrows we can turn Q into a triangulation, so the statement only needs to be proven for triangulations. By [19] we know that two trangulations of punctured surfaces can be turned into each other by a process of mutation.

This process removes an arrow to create a quadrangle and then puts in a new arrow. By lemma 4.6 this does not change $\mathtt{Tw\,Gtl}\,(\mathrm{RQ})$. □

5. THE RELATION WITH THE WRAPPED FUKAYA CATEGORY

A Liouville manifold is a manifold S with punctures equiped with a 1-form θ such that $\omega = d\theta$ is a symplectic form. The symplectic forms allows us to transform the 1-form into a vector field which generates a flow called the Liouville flow and we demand that the Liouville flow points towards the punctures near them. To any Liouville manifold one can associate an A_∞-category called the wrapped Fukaya category $\mathtt{Fuk}S$. We start with objects that are graded exact Lagrangian submanifolds which are invariant under the Liouville flow outside a compact subset of S.

The space of morphisms between two Lagrangians $\mathscr{L}_0$ and $\mathscr{L}_1$ is defined as the vector space spanned by the intersection points between $\mathscr{L}_1$ and the time 1 flow of $\mathscr{L}_0$ for a Hamiltonian $H : S \to \mathbb{R}$ which is quadratic at infinity.

$$\mathtt{Hom}(\mathscr{L}_0, \mathscr{L}_1) = \mathbb{C}^{\phi^1(\mathscr{L}_0) \cap \mathscr{L}_1} \text{ where } dH = \omega(\frac{d}{dt}\phi^t, -)$$

Note that such intersection points are also in one to one correspondences with time one chords between $\mathscr{L}_0$ and $\mathscr{L}_1$. These are curves $\gamma : [0,1] \to S : t \mapsto \phi^t(\gamma(0))$ such that $\gamma(0) \in \mathscr{L}_0$ and $\gamma(1) \in \mathscr{L}_1$.

Consider a sequence of intersection points $\mathscr{L}_k \xleftarrow{p_k} \mathscr{L}_{k-1} \cdots \mathscr{L}_1 \xleftarrow{p_1} \mathscr{L}_0$. The product is defined as

$$\mu_n(p_k, \dots, p_1) = \sum_{\mathscr{L}_0 \xleftarrow{q} \mathscr{L}_k} m_q q.$$

where m_g counts the signed number of immersed convex $k+1$-gons such that the edges map to $\phi^k\mathscr{L}_0, \phi^{k-1}\mathscr{L}_1, \dots \phi^1\mathscr{L}_{k-1}, \mathscr{L}_k$ and vertices to $\phi^{k-1}p_1, \phi^{k-2}p_2, \dots, p_k$ and q'. Here q' is the intersection point between $\phi^k\mathscr{L}_0$ and $\mathscr{L}_k$ such that q' gradually turns into q

if we look at the intersection points between $\phi^t\mathscr{L}_0$ and $\mathscr{L}_k$ and vary t from k to 1. Finally, to get the full wrapped Fukaya category we take the twisted completion[3]. The definition of the wrapped Fukaya category has a lot of technical issues for which we refer to [39] and [2].

In the case of a punctured surface, such Lagrangians can be seen as curves connecting the punctures. By tweaking the Liouville structure on the punctured surface we can ensure that each arrow a_i of an embedded quiver is an exact Lagrangian $\mathcal{L}_i$ and the punctures of the surface match the vertices of the quiver.

The Hamiltonian will make the flow spiral around the punctures, as illustrated below.





This implies that for every natural number n there is a time one chords between the two Lagrangians that circle around the puncture counterclockwise n times. Likewise in the algebra $\mathtt{Gtl}\,(\mathrm{RQ})$, there is also exactly one path between the two vertices that circle around the puncture n times. This means that there is a bijection between $\mathtt{Hom}_{\mathtt{Fuk}}(\mathscr{L}_i, \mathscr{L}_j)$ and $v_{a_i}\mathtt{Gtl}\,(\mathrm{RQ})v_{a_j}$. So, considered as a vector space the full subcategory of $\mathtt{F}_{\mathrm{Q}} \subset \mathtt{Fuk}S \setminus \mathrm{Q}_0$ containing the objects $\mathscr{L}_i$ and the category $\mathtt{Gtl}\,(\mathrm{RQ})$ are isomorphic.

Theorem 5.1. *Consider a quiver Q that splits a surface S and put a Liouville structure on $S \setminus Q_0$.*

1. *The full subcategory $\mathtt{fuk}(\mathrm{Q})$ of the wrapped Fukaya category $\mathtt{Fuk}S \setminus Q_0$ with objects the arrows of Q is $\mathtt{A}_\infty$-isomorphic to $\mathtt{Gtl}\,(\mathrm{RQ})$ with its generic $\mathtt{A}_\infty$-structure.*
2. *$\mathtt{Fuk}S \setminus Q_0$ is equivalent to $\mathtt{Tw}\,\mathtt{Gtl}\,(\mathrm{RQ}), \mu$.*

Proof. (Sketch) We already know that there is a linear isomorphism, so we only need to check the ordinary multiplications. The pictures of the compositions are easier to parse if we go to the universal cover of a closed neighborhood of a puncture. As such a neighborhood is a cylinder, the universal cover will be the upper half plane. The original Lagrangians will be vertical half lines and the flowed lines can be seen as sloped curves.


$\mathscr{L}_1$
α
$\mathbf{1}_{\mathscr{L}_0}$
$\mathscr{L}_0$ $\phi^1\mathscr{L}_0$



We can now draw a picture for the ordinary multiplication and the product of all arrows around a positive cycle in $\mathrm{R}Q$. The immersed polygon is shaded in both pictures.

---

[3] In principle we need to take the Karoubi completion, which ensures that direct summands of objects will also be objects.



$\Longrightarrow \mu(\beta,\alpha) = \beta\alpha$



$\Longrightarrow \mu(\gamma,\beta,\alpha) = \pm\mathbf{1}_{\mathscr{L}_0}$

Finally all products of shorter or longer sequences of arrows around a positive cycle are zero because there is no appropriate intersection point or there is a convexity problem. This establishes the first part of the theorem.

The second part is basically a consequence of the fact that $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ})$ does not depend on the quiver. Given any Lagrangian in $\mathtt{Fuk}S \setminus \mathrm{Q}_0$ we can add Lagrangians until we get a quiver Q that splits the surface. This implies that this particular Lagrangian sits in $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ})$. So every Lagrangian sits in a $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ})$ and $\mathtt{H\,Tw\,Gtl}\,(\mathrm{RQ}) \subset \mathtt{Fuk}(S \setminus \mathrm{Q}_0) \subset \mathtt{H\,Tw}\,(\mathtt{Gtl}\,\mathrm{RQ})$. □

6. CONSISTENT DIMER MODELS AND MATRIX FACTORIZATIONS

6.1. **Matrix factorizations in general.** Consider either an algebra $\mathtt{A}$ or a smooth algebraic variety X. A potential is defined as a central element $W \in Z(A)$ in the former case or a polynomial function $X \to \mathbb{C}$. The pair (X, W) is called a *commutative Landau-Ginzburg model*, the pair $(\mathtt{A}, W)$ is a *noncommutative Landau-Ginzburg model*.

A *matrix factorization* of a Landau-Ginzburg model is a diagram $\bar{P}$

$$\begin{tikzcd} P_0 \arrow[r, shift left, "p_0"] & P_1 \arrow[l, shift left, "p_1"] \end{tikzcd}$$

where P_i are projective $\mathtt{A}$-modules or vector bundles over X such that that $p_0p_1 = W$ and $p_1p_0 = W$.

Given 2 matrix factorizations $\bar{P}$ and $\bar{Q}$ we define $\mathtt{Hom}(\bar{P}, \bar{Q})$ as the following $\mathbb{Z}_2$-graded space of module morphisms/sheaf morphisms

$$\mathtt{Hom}(\bar{P}, \bar{Q}) = \underbrace{\mathtt{Hom}_{\mathtt{C}}(P_0, Q_0) \oplus \mathtt{Hom}_{\mathtt{C}}(P_1, Q_1)}_{\text{even}} \oplus \underbrace{\mathtt{Hom}_{\mathtt{C}}(P_0, Q_1) \oplus \mathtt{Hom}_{\mathtt{C}}(P_1, Q_0)}_{\text{odd}}.$$

On this space we have a differential of odd degree:

$$d\begin{pmatrix} f_{00} & f_{01} \\ f_{10} & f_{11} \end{pmatrix} := \begin{pmatrix} 0 & p_0 \\ p_1 & 0 \end{pmatrix}\begin{pmatrix} f_{00} & f_{01} \\ f_{10} & f_{11} \end{pmatrix} - \begin{pmatrix} f_{00} & -f_{01} \\ -f_{10} & f_{11} \end{pmatrix}\begin{pmatrix} 0 & q_0 \\ q_1 & 0 \end{pmatrix}.$$

It is easy to check that $d^2 = 0$.

The category of all matrix factorizations is denoted by $\mathtt{MF}(\mathtt{A}, W)$ or $\mathtt{MF}(X, W)$ and it has the structure of a $\mathbb{Z}_2$-graded dg-category or $\mathtt{A}_\infty$-category. In many cases it is interesting to look at small subcategories of this $\mathtt{A}_\infty$-category.

6.2. **The $\mathtt{A}_\infty$-algebra for a consistent dimer model.** Let Q be a zigzag consistent dimer model on a surface (with nonzero genus) and A its Jacobi algebra. The Jacobi algebra has a central element coming from the cycles

$$\ell := \sum_{v \in \mathrm{Q}_0} c_v$$

where $c_v \in \mathrm{Q}_2$ is a cycle starting in v. Note that the relations in the Jacobi algebra ensure that this expression does not depend on the choices of v and is indeed central.

For each arrow a we can define a matrix factorization of ℓ

$$\bar{P}_a := \begin{tikzcd} Ah(a) \arrow[r, shift left, "a"] & At(a) \arrow[l, shift left, "\bar{a}"] \end{tikzcd}$$

where $\bar{a}$ is defined such that $a\bar{a} \in \mathrm{Q}_2$.

We define $\mathtt{mf}(\mathrm{Q})$ as the dg-category with as objects the $\bar{P}_a$ and morphisms as in the previous section.

If $\tilde{\mathrm{Q}}$ is the universal cover of Q and π the cover map, then we can also define matrix factorizations in the universal cover $\bar{P}_{\tilde{a}}$ for every $\tilde{a} \in \tilde{\mathrm{Q}}_1$. By lifting paths to the universal cover we can easily see that

$$\mathtt{Hom}(\bar{P}_a, \bar{P}_b) = \bigoplus_{\tilde{b}, \pi(\tilde{b})=b} \mathtt{Hom}(\bar{P}_{\tilde{a}}, \bar{P}_{\tilde{b}}).$$

where we have chosen a fixed lift of a and vary over the lifts of b. This identification is also compatible with the differentials on both sides.

As we already know $\mathtt{mf}(\mathrm{Q})$ is a dg-algebra and hence its homology $\mathtt{Hmf}(\mathrm{Q})$ allows an $\mathtt{A}_\infty$-structure. Because the original dg-structure is compatible with the lift to the universal cover, this $\mathtt{A}_\infty$-structure will also be compatible.

6.3. **A nice basis for $\mathtt{Hmf}(\tilde{\mathrm{Q}})$.** Recall that the *zig ray* $\mathcal{Z}^+_{a_0}$ (zag ray $\mathcal{Z}^-_{a_0}$) of an arrow a_0 is the infinite sequence $\dots a_2 a_1 a_0$ of arrows in the universal cover, $\tilde{\mathrm{Q}}$, such that $a_{i+1}a_i$ sits in a positive cycle if i is even (odd) and in a negative cycle if i is odd (even). Given a piece of a zag ray $\mathcal{Z} = a_u \dots a_1 a_0$ with even length, we define the left (right) opposite path as the path formed by all arrows that are not in $\mathcal{Z}$ but are in positive (negative) cycles which meet $\mathcal{Z}$. If $\mathcal{Z}$ has odd length we take the left opposite path $a_{u-1} \dots a_0$ and the right opposite path of $a_u \dots a_1$.

We can put these two opposite paths in the correct entries of a 2×2-matrix and add an appropriate minus sign on the upper right entry.



gives $\zeta(\mathcal{Z}) = \begin{pmatrix} 0 & -\mathrm{opp}_1 \\ \mathrm{opp}_2 & 0 \end{pmatrix}$



gives $\zeta(\mathcal{Z}) = \begin{pmatrix} \mathrm{opp}_2 & 0 \\ 0 & \mathrm{opp}_1 \end{pmatrix}$

In this way we obtain an element in $\zeta(\mathcal{Z}) \in \mathtt{Hom}(P_{a_u}, P_{a_0})$ which is an off-diagonal or diagonal matrix, depending on whether the length of $\mathcal{Z}$ is even or odd.

Lemma 6.1. *If* Q *is a consistent dimer model on a surface with nonpositive Euler characteristic and* $\mathcal{Z}$ *is a zigzag path then both opposite paths are minimal (i.e. not a multiple of* ℓ*).*

Proof. From [12] we know there is a unique minimal path between every two vertices in the universal cover and such that every other path between these vertices is a power of ℓ times this path.

Let p be the minimal path in the universal cover that connects $h(\mathcal{Z})$ to $t(\mathcal{Z})$ and let a be the last arrow of $\mathcal{Z}$. First suppose that p does not cross $\mathcal{Z}$ and assume we cannot apply any reductions to p that make the piece that p and $\mathcal{Z}$ cut out smaller. We will prove the lemma by induction on the number of faces inside this piece.

If this number is 1 we are done because then the path is precisely the opposite path. If the number is bigger than 1, then the other zigzag path $\mathcal{Z}'$ that goes through the penultimate arrow of $\mathcal{Z}$ must cut the piece in two. Let b be the arrow where $\mathcal{Z}'$ intersects p and denote by p' the path which consists of the piece of p before b. If p is minimal then so must $p'a$ be and by induction it must be the path that goes opposite to $\mathcal{Z}'$.

The arrow b makes that bp' and a forteriori p can be reduced.



If p does cross $\mathcal{Z}$ in an arrow or a vertex, we can apply the above argument for each of the pieces separately and get the same result. ☐

Lemma 6.2. *The $\zeta(\mathcal{Z})$ for which $\mathcal{Z}$ starts in a and ends in b form a basis for* $\mathtt{H\,Hom}(\bar{P}_b, \bar{P}_a)$.

Proof. Because of $\mathbb{Z}_2$-degree and path-degree reasons we can look for basis elements of the forms

$$\begin{pmatrix} f_{00} & 0 \\ 0 & f_{11} \end{pmatrix}, \begin{pmatrix} 0 & f_{01} \\ f_{10} & 0 \end{pmatrix}$$

where the f_{ij} are paths up to a sign. We will only treat the first case. Being in the kernel of d implies that $f_{11} = a^{-1}(f_{00}b) = (\bar{a}f_{00})\bar{b}^{-1} \in \mathtt{Jac}\,(\mathrm{Q})$. These two conditions are the same as asking that $\bar{a}f_{00}b$ is a multiple of ℓ. Being in the image of d implies that f_{00} is a left multiple of a or a right multiple of $\bar{b}$ (which is equivalent to f_{11} being a right multiple of b or a left multiple of $\bar{a}$).

So we are looking for paths f_{00} such that $\bar{a}f_{00}b$ is a multiple of ℓ but neither $\bar{a}f_{00}$ or $f_{00}b$ is a multiple of ℓ. In particular this means that f_{00} and f_{11} are both minimal paths.

Now look at the paths f_{00} and f_{11} in the universal cover. If these 2 paths intersect at a vertex v we can split f_{ii} in two $f_{iv}vf_{vi}$. Now $\bar{a}f_{0v}$ and f_{1v} are both minimal and run between the same vertices so they must be the same. But then $f_{11} = a\bar{a}f_{0v}f_{v1}$ is a left multiple of $\bar{a}$ which is impossible. So a, b, f_{00} and f_{11} bound a simply connected piece S in the universal cover. After applying the relations we can assume that this piece is as small as possible.

Look at the zigzag path $\mathcal{Z}$ that starts from a and enters this simply connected piece. This zigzag path must leave the piece at an arrow c.



If c is in f_{00} then we can split $f_{00} = f_{0c}cf_{c0}$. Because of minimality f_{0c} and the opposite path running along the zigzag path from $h(a)$ to $h(c)$ must be equal. But then $f_{0c}c$ ends in $\bar{d}$ where d precedes c in the zigzag path. This contradicts the fact that we chose S as small as possible. Similarly $\mathcal{Z}$ cannot leave S through an arrow of f_{11}, so b must lie on the zigzag path through a. In order for $\mathcal{Z}$ to leave S at b, we must have that $\mathcal{Z}[i] = b$ with b even. ☐

Corollary 6.3. *In the universal cover $\tilde{\mathrm{Q}}$ the space* $\mathtt{H\,Hom}(\bar{P}_{\tilde{b}}, \bar{P}_{\tilde{a}})$ *is either zero dimensional or one dimensional, depending on whether $\tilde{b}$ sits in a zigzag ray from $\tilde{a}$ or not.*

Now we are going to express the A_∞-products in terms of these bases. To do this we will expand this basis to a full basis of $\mathtt{Hom}(\bar{P}_b, \bar{P}_a)$.

Consider the set of matrices

- $\begin{pmatrix} f_{00} & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & f_{01} \\ 0 & 0 \end{pmatrix}$ where f_{00} and f_{01} are paths.

- $\begin{pmatrix}0 & 0\\ 0 & f_{11}\end{pmatrix}, \begin{pmatrix}0 & 0\\ f_{10} & 0\end{pmatrix}$ where f_{11} and f_{10} are paths and $af_{11}b^{-1}, af_{10}\bar{b}^{-1} \not\in \mathtt{Jac}\,(\mathrm{Q})$

and let U be the span of these matrices.

By construction it is clear that the matrices above are linearly independent and

$$\mathtt{Hom}(P_a, P_b) = U \oplus \mathtt{H}\,\mathtt{Hom}(P_a, P_b) \oplus d(U).$$

This split allows us to define a codifferential

$$h : \mathtt{Hom}(P_a, P_b) \to \mathtt{Hom}(P_a, P_b) : u_1 \oplus \zeta(\mathcal{Z}_a^{\pm,j}) \oplus du_2 \mapsto u_2 \oplus 0 \oplus 0$$

Lemma 6.4.

- *$dhd = d$,*
- *$h^2 = 0$,*
- *$h(\zeta(\mathcal{Z})) = 0$.*
- *h respects the split $\mathtt{Hom}(\bar{P}_a, \bar{P}_b) = \bigoplus_{\tilde{b}, \pi(\tilde{b})=b} \mathtt{Hom}(\bar{P}_{\tilde{a}}, \bar{P}_{\tilde{b}})$.*

Proof. These facts follow easily from the construction. □

We are now going to determine the A_∞-structure on $\mathtt{H}\,\mathtt{mf}(\mathrm{Q})$. First we start with the ordinary multiplication.

Theorem 6.5. *Suppose $\mathcal{Z}_1$ is a zigzag path from b to a and $\mathcal{Z}_2$ is a zigzag path from a to c*

$$\mu(\zeta(\mathcal{Z}_1), \zeta(\mathcal{Z}_2)) = \begin{cases}\zeta(\mathcal{Z}_2 a^{-1}\mathcal{Z}_1) & \textit{if } \mathcal{Z}_2 a^{-1}\mathcal{Z}_1 \textit{ is a zigzag path}\\ 0 & \textit{otherwise}\end{cases}$$

Proof. From the formula for the $\mathtt{A}_\infty$-structure we get

$$\mu(\zeta(\mathcal{Z}_1), \zeta(\mathcal{Z}_2)) = (1 - hd + dh)(\zeta(\mathcal{Z}_1)\zeta(\mathcal{Z}_2))$$

Now $\zeta(\mathcal{Z}_1)\zeta(\mathcal{Z}_2)$ is a matrix that either consists of 2 paths (possibly with signs) on the off-diagonals or on the diagonals. The expression $(1 - hd + dh)(\zeta(\mathcal{Z}_1)\zeta(\mathcal{Z}_2))$ will be nonzero only if these paths are opposite paths of some other zigzag path.

If $\mathcal{Z}_2 a^{-1}\mathcal{Z}_1$ is a zigzag path, one can easily check that $\zeta(\mathcal{Z}_1)\zeta(\mathcal{Z}_2) = \zeta(\mathcal{Z}_2 a^{-1}\mathcal{Z}_1)$ and hence $\mu(\zeta(\mathcal{Z}_1), \zeta(\mathcal{Z}_2)) = \zeta(\mathcal{Z}_2 a^{-1}\mathcal{Z}_1)$.

If $\mathcal{Z}_1$ and $\mathcal{Z}_2$ are zigzag paths in different directions, we will show that $\mathtt{H}\,\mathtt{Hom}(\bar{P}_b, \bar{P}_c) = 0$ in the universal cover. If this were not the case then there must be a zigzag path $\mathcal{Z}$ from c to b.

If $\mathcal{Z}_2$ lies to the left of $\mathcal{Z}_1$ then in order for $\mathcal{Z}$ to intersect with $\mathcal{Z}_2$, $\mathcal{Z}$ must also lie to the left of $\mathcal{Z}_1$ and therefore the length of $\mathcal{Z}_1$ must be odd.



For similar reasons the length of $\mathcal{Z}_2$ must be odd while the length of $\mathcal{Z}$ must be even. This is impossible because of the degree of $\mu(\zeta(\mathcal{Z}_1), \zeta(\mathcal{Z}_2))$ would then be even while the degree of $\zeta(\mathcal{Z})$ is odd. □

Theorem 6.6. *Let $a_1 \dots a_k$ be a positive or negative cycle in Q then*

$$\mu(\zeta(a_1 a_2), \zeta(a_2 a_3), \dots, \zeta(a_l a_{l+1})) = \begin{cases}0 & l \neq k\\ \zeta(a_1) & l = k\end{cases}$$

Proof. First note that

$$\zeta(a_ia_{i+1}) = \begin{pmatrix} 0 & -t(a_i) \\ a_{i+2}\dots a_{i+k-1} & 0 \end{pmatrix}.$$

Therefore

$$h(\zeta(a_ia_{i+1})\zeta(a_{i+1}a_{i+2})) = \begin{pmatrix} 0 & 0 \\ -a_{i+3}\dots a_{i+k-1} & 0 \end{pmatrix}$$

and by induction

$$\begin{aligned} &h(\dots h(h(\zeta(a_ia_{i+1})\zeta(a_{i+1}a_{i+2})\zeta(a_{i+2}a_{i+3})\dots)\zeta(a_{i+r}a_{i+r+1}) \\ &= \begin{pmatrix} 0 & 0 \\ -a_{i+r+2}\dots a_{i+k-1} & 0 \end{pmatrix}. \end{aligned}$$

if $r+2 \leq k-1$ and zero otherwise.

If we multiply two such expressions together we get zero. If we multiply one such expression on the left with $\zeta(a_{i-1}a_i)$ we get a matrix with only the upper left diagonal nonzero. If we apply h to this we also get zero.

The only trees that might give a nonzero contribution to μ are



$$\pi(h(\dots h(h(\zeta(a_1a_2)\zeta(a_2a_3)\zeta(a_{i+2}a_{i+3})\dots)\zeta(a_ra_{r+1}))$$



$$\pi(\zeta(a_1a_2)h(\dots(h(\zeta(a_2a_3)\zeta(a_{i+2}a_{i+3})\dots)\zeta(a_ra_{r+1}))$$

Both expressions can only give a nonzero result if $r = k$. Indeed by degree reasons these expressions are even: $r\deg\zeta(a_1a_2) + (r-2)\deg h + \deg\pi = 0$, but the only path that goes opposite a zigzag path of odd length which is also a subpath of $a_1\dots a_{i+r}$ is the trivial path and this happens when $r = 0 \mod k$. If $r > k+1$ the expression $h(h\dots h(\dots))$ will be zero.

If $r = k$ then we see that

$$\begin{aligned} &h(\dots h(h(\zeta(a_1a_2)\zeta(a_2a_3)\zeta(a_{i+2}a_{i+3})\dots)\zeta(a_ra_{r+1})) \\ &+\zeta(a_1a_2)h(\dots(h(\zeta(a_2a_3)\zeta(a_{i+2}a_{i+3})\dots)\zeta(a_ra_{r+1})) \\ &= \begin{pmatrix} h(a_1) & 0 \\ 0 & t(a_1) \end{pmatrix} = \zeta(a_1). \end{aligned}$$

Because the latter is a basis element of the homology, $\pi(\zeta(a_1)) = \zeta(a_1)$. □

7. DIMER DUALITY AS MIRROR SYMMETRY

Let Q be any, not necessarily consistent, dimer. We define its mirror dimer $\reflectbox{Q}$ as follows

1. The vertices of $\reflectbox{Q}$ are the zigzag cycles of Q.
2. The arrows of $\reflectbox{Q}$ are the arrows of Q, $h(a)$ is the zigzag cycle coming from the zig ray, and $t(a)$ is the cycle coming from the zag ray.
3. The positive faces of $\reflectbox{Q}$ are the positive faces of Q.
4. The negative faces of $\reflectbox{Q}$ are the negative faces of Q in reverse order.

We illustrate this with a couple of examples:



Remark 7.1. The dual can also be obtained by cutting the dimer along the arrows, flipping over the clockwise faces, reversing their arrows and gluing everything back again. This construction is basically the same construction that was introduced by Feng, He, Kennaway and Vafa in [13] applied to all possible dimers.

Lemma 7.2. *$\mathrm{Q} \mapsto \reflectbox{Q}$ is an involution on the set of all dimer models.*

Proof. Let $\mathcal{Z} = a_1 \dots a_k$ be a zigzag cycle in $\reflectbox{$\mathrm{Q}$}$. If i is odd then $a_i a_{i+1}$ sits in a positive cycle in $\reflectbox{$\mathrm{Q}$}$ while $a_i a_{i+1}$ sits in a negative cycle of $\reflectbox{$\mathrm{Q}$}$ when i is even.

This implies that $a_i a_{i+1}$ sits in a positive cycle in Q for i odd while $a_{i+1} a_i$ sits in a negative cycle of Q when i is even. Hence in Q the odd arrows of $\mathcal{Z}$ must have the same head in Q which is equal to the tail of the even arrows. This means there is a well defined map from the zigzag paths in $\reflectbox{$\mathrm{Q}$}$ and the vertices of Q. Because a is an even arrow in its zig cycle and an odd in its zag cycle, under this map the zig cycle and the zag cycle of a in $\reflectbox{$\mathrm{Q}$}$ correspond to the original head and tail. Finally this map is a bijection because given any vertex $v \in \mathrm{Q}$ we can make a zigzag cycle in $\reflectbox{$\mathrm{Q}$}$ by listing all arrows incident with v in a clockwise direction. □

Theorem 7.3. *If Q is a consistent dimer then $\mathtt{Hmf}(\mathrm{Q})$ is $\mathtt{A}_\infty$-isomorphic to $\mathtt{Gtl}\,(\mathrm{R}\reflectbox{Q}), \mu$.*

Proof. The underlying category of $\mathtt{Hmf}(\mathrm{Q})$ is indeed isomorphic to $\mathtt{Gtl}\,(\mathrm{R}\reflectbox{Q})$. The paths in $\mathtt{Gtl}\,(\mathrm{R}\reflectbox{Q})$ are those that cycle around vertices of $\reflectbox{$\mathrm{Q}$}$, which are precisely the zigzag paths of Q. Two paths multiply to zero in $\mathtt{Gtl}\,(\mathrm{R}\reflectbox{Q})$ if they go around different vertices in $\reflectbox{$\mathrm{Q}$}$ just as the product of $\zeta(p_1) \cdot \zeta(p_2)$ is zero if they belong to different zigzag cycles. Therefore we clearly have an isomorphism of categories.

Finally the $\mathtt{A}_\infty$-structure is $\mathtt{A}_\infty$-isomorphic to μ because of lemma 6.6 and theorem 4.4. □

Corollary 7.4. *For any consistent dimer Q we have that $\mathtt{Hmf}(\mathrm{Q})$ and $\mathtt{Fuk}(|\reflectbox{Q}| \setminus \reflectbox{Q}_0)$ are $\mathtt{A}_\infty$-isomorphic $\mathtt{A}_\infty$-categories.*

Note that the above result applies to all consistent dimers, not only those on the torus. This means that the Jacobi algebra of these dimers is not necessarily Noetherian. So in some cases this result does not derive from a commutative instance of mirror symmetry in the background. However if one restricts to dimer models on a torus it is possible to recover the commutative picture from the dimer itself.

8. RECOVERING THE COMMUTATIVE PICTURE

Let Q be a consistent dimer model on a torus and let $\mathtt{A} = \mathtt{Jac}\,(\mathrm{Q})$ be its Jacobi algebra. Fix a vertex o which we will call the trivial vertex. Now choose 2 cyclic paths x, y through o that are not multiples of ℓ and form a basis for the homology of the torus. Finally let z be any positive cycle through o.

A *perfect matching* $\mathcal{P}$ is a subset of the arrows such that every positive or negative cycle contains exactly one arrow of $\mathcal{P}$. Every perfect matching defines a degree function on A

by setting the degree of an arrow equal to 1 if it is in the matching and zero otherwise. A collection of perfect matchings $\{\mathcal{P}_1, \dots, \mathcal{P}_k\}$ is called *θ-stable*[4] if there is a path from o to every vertex that has degree zero for all the matchings in the collections.

For every stable collection $S = \{\mathcal{P}_1, \dots, \mathcal{P}_k\}$ we define a cone in $N = \mathbb{R}^3$

$$\sigma_S = \mathbb{R}^+(\mathcal{P}_1(x), \mathcal{P}_1(y), \mathcal{P}_1(z)) + \cdots + \mathbb{R}^+(\mathcal{P}_k(x), \mathcal{P}_k(y), \mathcal{P}_k(z))$$

Theorem 8.1 (Ishii-Ueda-Mozgovoy)**.** *The collection of cones σ_S where S is a stable set of matchings forms a fan and the toric variety of this fan, $\tilde{X}$ is a crepant resolution of $X = \mathsf{Spec}Z(A)$.*

If we intersect the fan with the plane at height $z = 1$, we get a convex polygon, which is subdivided in triangles and on each integral point of the polygon sits a unique perfect matching. These lattice points will also give us a basis for the toric divisors of $\tilde{X}$. So any line bundle over $\tilde{X}$ can be identified with a $\mathbb{Z}$-linear combination of stable perfect matchings. In this language we have the following:

Theorem 8.2 (Ishii-Ueda [22]-Mozgovoy-Bender[33])**.** *There is an equivalence of categories $\mathscr{F} : \mathtt{D}^b\mathtt{Mod}A \to \mathtt{D}^b\mathtt{Coh}\,\tilde{X}$, which maps the projective $v\mathtt{A}$ to the line bundle $\sum \mathcal{P}_i(p_v)\mathcal{P}_i$ where p_v is any path from o to v and the sum runs over all stable matchings.*

In [23] Ishii and Ueda use this result to get an equivalence of categories between the corresponding categories of matrix factorizations.

Theorem 8.3 (Ishii-Ueda-[23])**.** *There is an equivalence of categories $\mathscr{F} : \mathtt{H}^0\mathtt{MF}(A, \ell) \to \mathtt{H}^0\mathtt{MF}(\tilde{X}, \ell)$.*

The last theorem implies that any of the matrix factorizations $\bar{P}_a$ is transformed into a matrix factorization of line bundles $\bar{\mathscr{L}}_a$ over $\tilde{X}$.

We illustrate the theorems with one particular dimer model. We take vertex 1 to be the trivial one.

Dimer



Stable Matchings

$\mathcal{P}_1 = \{b, w\}$
$\mathcal{P}_2 = \{d, z\}$
$\mathcal{P}_3 = \{a, y\}$
$\mathcal{P}_4 = \{c, x\}$
$\mathcal{P}_5 = \{a, b\}$

Fan



Linebundles

$\mathscr{L}_1 = \mathcal{O}$
$\mathscr{L}_2 = \mathcal{O}_{V_2}$
$\mathscr{L}_3 = \mathcal{O}_{V_1+V_2}$
$\mathscr{L}_4 = \mathcal{O}_{V_1+V_2+V_4}$

The only thing that is missing to get an equivalence

$$\mathtt{H}^0\mathtt{MF}(\tilde{X}, \ell) \cong \mathtt{H}^0\mathtt{Fuk}|\mathcal{Q}| \setminus \mathcal{Q}_0$$

is a proof that the objects $\bar{\mathscr{L}}_a$ generate $\mathtt{H}^0\mathtt{MF}(\tilde{X}, \ell)$. We will briefly sketch a proof of this. In [36] Orlov proved that we have an equivalence of categories

$$\mathtt{H}^0\mathtt{MF}(\tilde{X}, \ell) \cong \frac{\mathtt{D}^b\mathtt{Coh}\,(\tilde{X}_0)}{\mathtt{Perf}(\tilde{X}_0)} \text{ with } \tilde{X}_0 := \ell^{-1}(0).$$

The second category is the quotient of the derived category of the zero fiber of ℓ by its perfect complexes. This category is also called the category of singularities of $\tilde{X}_0$. The functor that realizes this equivalences maps a matrix factorization $P_0 \underset{p_1}{\overset{p_0}{\rightleftarrows}} P_1$. to $\mathtt{Cok}p_0$. If we can prove that $\mathtt{D}^b\mathtt{Coh}\,(\ell^{-1}(0))$ is generated by the $\mathtt{Cok}\mathscr{F}a$ for $a \in \mathrm{Q}_1$ we are done.

For each cone σ of co-dimension i in the fan we get a dimension i closed subvariety Y_σ of $\tilde{X}_0$ which is the intersection of the toric invariant divisors that generate the cone. If we can find for each σ an object $\mathscr{M}_\sigma$ that is supported on an open subvariety of Y_σ, then the collection $\mathscr{M}_\sigma$ will generate $\mathtt{D}^b\mathtt{Coh}\,(\tilde{X}_0)$. This is true because $\mathtt{D}^b\mathtt{Coh}\,(\tilde{X}_0)$ is generated by the subcategories $\mathtt{D}^b_{Y_\sigma}\mathtt{Coh}\,\tilde{X}_0$, containing the objects with homology supported on the Y_σ. In fact one only needs the Y_σ with codimension zero because these are the connected components of $\tilde{X}_0$. Each of the $Y_\sigma \setminus \cup_{\tau\neq\sigma} Y_\tau$ is affine and hence $\mathscr{M}_i$ together with the subcategories $\mathtt{D}^b\mathtt{Coh}_{\,Y_\sigma\cap Y_\tau}\tilde{X}_0$ generate the whole $\mathtt{D}^b\mathtt{Coh}_{\,Y_\sigma}(\tilde{X}_0)$.

---

[4] The θ we use is negative for the trivial and positive for all other vertices.

Let σ be a cone generated by $\{\mathcal{P}_{i_1},\dots,\mathcal{P}_{i_k}\}$. Consider $\mathscr{U}_\sigma = \bigoplus_{a\in\cup\mathcal{P}_{i_j}} \mathtt{Cok}a$. This is a sheaf over $\tilde{X}_0$ which has maximal rank at the locus where all $a \in \cup\mathcal{P}_{i_j}$ are zero. This is precisely Y_σ and hence $\mathcal{U}_\sigma$ generates a sheaf $\mathscr{M}_\sigma$ which has rank one on an open subspace of Y_i and is zero elsewhere.

All this allows us to conclude:

Theorem 8.4. *Given a consistent dimer model on a torus* Q*, and* $\tilde{X}$ *a crepant resolution of* $\mathsf{Spec}Z(\mathtt{Jac}\,\mathrm{Q})$ *we have the following equivalences.*

$$\mathtt{H}^0\mathtt{MF}(\tilde{X},\ell) \cong \mathtt{H}^0\mathtt{MF}(\mathtt{Jac}\,\mathrm{Q},\ell) \cong \mathtt{H}^0\mathtt{Fuk}|\wp|\setminus\wp_0$$

We end with a graphical connection between the surface of the mirror dimer and the polygon constructed from the original dimer.

Theorem 8.5. *Let* Q *be a consistent dimer on a torus. The mirror dimer* $\wp$ *gives us a surface* $|\wp|\setminus \wp_0$ *of genus* g *with* k *punctures, where* g *is the number of internal lattice points of the polygon associated to* Q *and* k *is the number of boundary lattice points.*

Proof. For every lattice point in the polygon there is precisely one stable perfect matching. If we take two consecutive stable perfect matchings $\mathcal{P}_1,\mathcal{P}_2$ on the boundary, then the arrows in $\mathcal{P}_1\cup\mathcal{P}_2\setminus\mathcal{P}_1\cap\mathcal{P}_2$ form a zigzag path and any zigzag path can be constructed in this way (for a proof of this see [**?**]). This implies there is a bijection that maps the line segments on the boundary between the lattice points to zigzag paths and hence the number of zigzag paths equals the number of boundary lattice points.

The equivalence between the K-groups of $\mathtt{D}^b\mathtt{Coh}\,\tilde{X}$ and $\mathtt{D}^b\mathtt{ModJac}\,(\mathrm{Q})$ shows that the number of maximal cones in the fan of $\tilde{X}$ equals the number of vertices in Q. The former equals twice the area of the polygon because the fan cuts the polygon in elementary triangles. By Pick's theorem this area is $B/2+I-1$ where B is the number of boundary lattice points and I the number of internal lattice points. From this discussion we get that

$$\#\,\wp_0 = \#\{\text{zigzag paths in Q}\} = B \text{ and } \#\mathrm{Q}_0 = B+2I-2.$$

The genus of $|\wp|$ is

$$\begin{aligned} g &= 1-\frac{\#\,\wp_0 - \#\,\wp_1 + \#\,\wp_2}{2}\\ &= 1-\frac{\#\mathrm{Q}_0-(2I+2)-\#\mathrm{Q}_1+\#\mathrm{Q}_2}{2}\\ &= 1-\frac{(2I+2)+0}{2} = I \end{aligned}$$

The first step follows because $\#\,\wp_i = \#\mathrm{Q}_i$ for $i=1,2$, while the second step is a consequence of the fact that Q sits in a torus. $\square$

Remark 8.6. This wellknown by physicists and can be found in [13]. Graphically we can say that the mirror surface $|\wp|\setminus\wp_0$ is a tubular neighborhood of the dual diagram of the fan associated to $\tilde{X}$.


Fan Dual $|\wp|\setminus\wp_0$

9. APPENDIX: HOCHSCHILD COHOMOLOGY AND $\mathtt{A}_\infty$-STRUCTURES

In this appendix we look at the connection between Hochschild cohomology and $\mathtt{A}_\infty$-structures and calculate the Hochschild cohomology of the gentle categories coming from rectified dimers.

Then we will define certain $\mathtt{A}_\infty$-structures on these gentle categories and use our calculation to find criteria for when a given $\mathtt{A}_\infty$-structure will be $\mathtt{A}_\infty$-isomorphic to one of our defined ones.

In this appendix we will assume that Q itself is a rectified dimer, so in particular every vertex has 2 arrows arriving and two arrows leaving.

9.1. **Hochschild cohomology and $\mathtt{A}_\infty$-structures.** The discussion in this section closely matches [7]. If we are interested in $\mathtt{A}_\infty$-structures on an ordinary category $\mathtt{B} = \mathtt{Gtl}\,\mathrm{Q}$ up to $\mathtt{A}_\infty$-isomorphism, we need to study the Hochschild cohomology of $\mathtt{B}$.

A length n multi-functor $\mathcal{F}$ consists of linear maps for each sequence of $n+1$ objects $X_0, \dots, X_n \in \mathtt{Ob}\,\mathtt{B}$.

$$\mathcal{F} : \mathtt{Hom}_\mathtt{B}(X_1, X_0) \otimes \cdots \otimes \mathtt{Hom}_\mathtt{B}(X_n, X_{n-1}) \to \mathtt{Hom}_\mathtt{C}(X_n, X_0).$$

The set of all length n-multifunctors forms a $\mathbb{Z}_2$-graded vector space which we denote by $\mathcal{M}^n$.

We can construct a differential $d : \mathcal{M}^n \to \mathcal{M}^{n+1}$

$$\begin{aligned} d\mathcal{F}(f_1, \dots, f_{n+1}) =& f_1\mathcal{F}(f_2, \dots, f_{n+1}) \\ -\mathcal{F}(f_1f_2, \dots, f_{n+1}) &+ \mathcal{F}(f_1, f_2f_3, \dots, f_{n+1}) \cdots \pm \mathcal{F}(f_1, \dots, f_{n-1}f_n) \\ &\mp \mathcal{F}(f_1, \dots, f_{n-1})f_n \end{aligned}$$

The *Hochschild cohomology* of $\mathtt{B}$ is defined as the cohomology of the complex $\mathcal{M}^\bullet$:

$$\mathtt{HochC}(\mathtt{B}) := \mathtt{H}\,(\mathcal{M}^\bullet).$$

The importance of the Hochschild cohomology of $\mathtt{C}$ can be seen by the following 2 lemmas.

Lemma 9.1. [7] *Let $\mu_i, i \le k$ be a sequence of multifunctors with $\mu_1 = 0$ and $\mu_2 = \cdot$ then we can rewrite $[\mathrm{M_k}]$ as*

$$d\mu_k = \Phi$$

where Φ is an expression calculated from the $\mu_i, i < k$ and $d\Phi = 0$ if all $[\mathrm{M_i}]$ for $< k$ hold.

Lemma 9.2. [7] *Let μ and ν be two $\mathtt{A}_\infty$-structures on $\mathtt{B}$ and let $\mathcal{F}_i$, $i \le k$ be a sequence of multifunctors with $\mathcal{F}_1 = \mathbf{1}$ then we can rewrite $[\mathrm{F_{k+1}}]$ as*

$$d\mathcal{F}_k = \Psi$$

where Ψ is an expression calculated from the μ_i, the ν_i and the $\mathcal{F}_i, i < k$. Moreover $d\Psi = 0$ if all $[\mathrm{F_i}]$ for $i < k$ hold.

These lemmas imply that we can find a solution for μ_k (or $\mathcal{F}_k$) if and only if the homology class of Φ (or Ψ) is trivial. So calculating the Hochschild cohomology of $\mathtt{B}$ is crucial if one wants to classify $\mathtt{A}_\infty$-structures on $\mathtt{B}$.

9.2. **Hochschild cohomology of the gentle categories.** Now we will calculate the Hochschild cohomology of the gentle category $\mathtt{B} = \mathtt{Gtl}\,(\mathrm{Q})$, which we will view as an algebra in this section. For this we will use a minimal bimodule resolution for $\mathtt{B}$, which can be obtained by a result by Bardzel [8].

Theorem 9.3 (Bardzel)**.** *Suppose $\mathtt{B}$ is the path algebra of a quiver Q modulo relations of length two. Let $Z_k \subset \mathbb{C}Q$ be the vector space spanned by paths of length k of which all subpaths of length 2 are zero in $\mathtt{B}$.*

The terms in minimal bimodule resolution $\mathcal{P}^\bullet$ are given by $\mathcal{P}_k = \mathtt{B} \otimes Z_k \otimes \mathtt{B}$ where the tensor product is taken over $\mathbb{C}Q_0$. The maps between the terms have the following form

$$1 \otimes b_1 \dots b_k \otimes 1 \mapsto b_1 \otimes b_2 \dots b_k \otimes 1 - (-1)^k \otimes b_1 \dots b_{k-1} \otimes b_k$$

As is well known the Hochschild cohomology of $\mathtt{B}$ is the cohomology of the complex

$$\mathtt{Hom}_{\mathtt{B}\otimes\mathtt{B}^{opp}}(\mathcal{P}^\bullet, \mathtt{B}).$$

Note that both this complex and $\mathcal{P}^\bullet$ are $\mathbb{Z}Q_1$-graded with degree zero differentials. Moreover in the case of these gentle algebras, the bimodule resolution also splits as a direct sum where each summand $\mathcal{P}^\bullet_c$ only contains the paths that go around one cycle $c \in \mathrm{Q}_2^+$. So let us focus on one such cycle $c = b_1 \dots b_l$ and set b_k to be $b_{k \mod l}$ for all $k \in \mathbb{Z}$.

Clearly

$$\mathtt{Hom}_{\mathtt{B}\otimes\mathtt{B}^{opp}}(\mathtt{B} \otimes b_i \dots b_j \otimes \mathtt{B}, \mathtt{B}) \cong h(b_i)\mathtt{B}t(b_j)$$

and we will use the notation $[b_i \dots b_j \to x]$ for the morphism that maps $b_i \dots b_j$ to $x \in h(b_i)Bt(b_j)$. By lemma 4.3, the space $h(b_i)\mathtt{B}t(b_j)$ is only nonzero in 3 cases.

1. If $|b_i \dots b_j| = -1 \mod l$ then $h(b_i)\mathtt{B}t(b_j) = b_{i-1}^{-1}z\mathbb{C}[z]$ where z is the unique negative cycle through b_{i-1}.
2. If $|b_i \dots b_j| = 0 \mod l$ then $h(b_i)\mathtt{B}t(b_j) = h(b_i)\mathbb{C}[z_1, z_2]/(z_1z_2)$ where z_1, z_2 are the two negative cycles that start in $h(b_i)$.
3. If $|b_i \dots b_j| = 1 \mod l$ then $h(b_i)\mathtt{B}t(b_j) = b_i\mathbb{C}[z]$ where z is the unique negative cycle through b_i.

With the notation above, the differential on $\mathtt{Hom}_{\mathtt{B}\otimes\mathtt{B}^{opp}}(\mathcal{P}^\bullet, \mathtt{B})$ becomes

$$d[b_i \dots b_j \to x] = [b_{i-1} \dots b_j \to b_{i-1}x] - (-1)^{j-i}[b_i \dots b_{j+1} \to xb_{j+1}]$$

1. If $|b_i \dots b_j| = j-i+1 = -1 \mod l$ then $d[b_i \dots b_j \to b_{i-1}^{-1}z^u] = [b_{i-1} \dots b_j \to z^u] - (-1)^{j-i}[b_i \dots b_{j+1} \to b_{i-1}^{-1}z^u b_{i+1}]$. Both terms only appear for the differential of $[b_i \dots b_j \to b_{i-1}^{-1}z^u]$, so d acts injectively on $\mathtt{Hom}(\mathcal{P}_c^{j-i+1}, \mathtt{B})$.
2. If $|b_i \dots b_j| = 0 \mod l$ and $x = z_1^u$ or $x = z_2^u$, $u > 0$ then one term of $d[b_i \dots b_j \to x]$ is zero and the other nonzero. Every such term appears exactly twice:

   $$\begin{cases} d[b_i \dots b_j \to x] = \pm d[b_{i-1} \dots b_{j-1} \to b_{i-1}xb_{j-1}^{-1}] & \text{if } b_{i-1} \text{ sits in } x \\ d[b_i \dots b_j \to x] = \pm d[b_{i+1} \dots b_{j+1} \to b_i^{-1}xb_i] & \text{if } b_i \text{ sits in } x \end{cases}$$

   The sum/difference gives us an element in the kernel which is in the image of d.

   For $x = h(b_i)$ we get that both terms are nonzero and it is easy to check that each term appears twice: once for $d[b_i \dots b_j \to h(b_i)]$ and once for $d[b_{i\pm1} \dots b_{j\pm1} \to h(b_{i\pm1})]$ and therefore the homology in degree $i - j + 1 = 0 \mod l$ is generated by elements of the form

   $$\sum_{0 \le u \le j-i} (-1)^{u(j-i)}[b_{i+u} \dots b_{j+u} \to h(b_{i+u})]$$
3. If $|b_i \dots b_j| = 1 \mod l$ then d acts as zero because for any $x = b_iz^u$ the terms of the differential each contain two consecutive arrows of c. As we have seen above, each term $[b_i \dots b_j \to b_iz^u]$ is in the image of d if $u > 0$. If $u = 0$ then $[b_i \dots b_j \to b_i] - (-1)^{j-i+1}[b_i \dots b_j \to b_j]$ is in the image of d, clearly this implies that

   $$\sum_{0 \le u \le j-i} (-1)^{u(j-i+1)}[b_{i+u} \dots b_{j+u} \to b_{i+u}]$$

   is the only element we need to add to the image to span the whole space.

From this discussion, we can conclude that

Proposition 9.4. *The homology of* $\mathtt{Hom}(\mathcal{P}^\bullet_c, \mathtt{B})$ *in degrees* $nl - 1$ *is zero, in degrees* nl *it is one and spanned by*

$$\Omega_0^{c,n} = \sum_{i=1}^{nl}(-1)^{i(nl-1)}[b_i \dots b_{i+nl-1} \to h(b_i)],$$

and in degrees $nl+1$ it is one and spanned by

$$\Omega_1^{c,n} = \sum_{i=1}^{nl} (-1)^{i(nl+1)} [b_i \dots b_{i+nl} \to b_i].$$

Now we go back to the original viewpoint of the Hochschild cohomology. Given a multifunctor Ψ of length k with $d\Psi = 0$, how can we detect whether its cohomology class is zero? The idea is to look at the image of k-tuples of the form $(b_1, \dots, b_k)$ where the b_i are consecutive arrows in a positive cycle:

Lemma 9.5. *Let Ψ be a multifunctor of length u with $d\Psi = 0$ then $\Psi \in Kerd$ if $\Psi(b_1, \dots, b_u)$ has no length 0 terms or length 1 terms for every $(b_1, \dots, b_u)$ where the b_i are consecutive arrows in a positive cycle.*

Proof. For $u = 0 \mod l$ the homology class $\Omega_0^{c,n}(b_i, \dots, b_{nl-1}) = h(b_i)$ contains a length 0 term. Now if Π is any multifunctor of length $nl-1$ then one can check that

$$d\Pi(b_i, \dots, b_{i+nl-1}) = b_i \Pi(b_{i+1}, \dots, b_{i+nl-1}) - 0 + \dots \mp 0 \pm \Pi(b_i, \dots, b_{i+nl-2}) b_{i+nl-1}$$

which has no terms of length 0 or 1 because neither $\Pi(b_{i+1}, \dots, b_{i+nl-1})$ nor $\Pi(b_i, \dots, b_{i+nl-2})$ can be a vertex.

Similarly for $u = 1 \mod l$, if Π is any multifunctor of length nl then we can write $d\Pi(b_i, \dots, b_{i+nl}) = \alpha_i b_i + \dots$ where $\alpha_i \in \mathbb{C}$. Now one easily checks that $\sum_{i=1}^{nl} (-1)^{(nl+1)i} \alpha_i = 0$. But if we replace $d\Pi$ by $\Omega_1^{c,n}$ we get that $\sum_{i=1}^{nl} (-1)^{(nl+1)i} \alpha_i = nl \neq 0$. □

Another ingredient we will need is an interesting collection of elements in Ker$d$. Choose for each arrow a and each $n \geq 1$ a number $\zeta_{a,n} \in \mathbb{C}$. Define the multifunctor g_k on $\mathtt{Gtl}\,(\mathrm{Q})$ by $g_k(x_1, \dots, x_k) = \zeta_{a,n} sat$ if $x_1 = sa$, $x_k = at$ and for $i \in [2, k-1]$ all x_i are arrows and $ax_2 \dots x_{k-1}$ is the n^{th} power of a positive cycle in Q. For other cases where the x_i are paths set g_k zero and extend it multilinearly.

Lemma 9.6. $dg_k = 0$.

Proof. Note that g_k is well defined because through every arrow there is only one positive cycle, so sat is uniquely determined by $x_1, \dots, x_k$.

Let $u_1, \dots, u_{k+1}$ be paths and suppose $dg_k(u_1, \dots u_{k+1}) \neq 0$, then there must at least be one term in the expression that is nonzero.

- If $u_1 g_k(u_2, \dots, u_{k+1}) \neq 0$ then $u_2 = sa, u_{k+1} = at$ for some a and $u_1 sat \neq 0$. The second term is $-g_k(u_1u_2, \dots, u_{k+1}) = -\zeta_{n,a} u_1 sat$. The last term is nonzero because u_k is an arrow b such that a and b sit in a positive cycle and hence $ba = 0$ but if $g_k(u_1, u_2, \dots, u_k)u_{k+1} \neq 0$ then $u_1 = s'b$ which contradicts $u_1 sat \neq 0$. Finally all intermediate terms are also zero because they contain the product of two consecutive arrows in a positive cycle.
- If $g_k(u_1u_2, \dots, u_{k+1}) \neq 0$ then $u_1u_2 = sa$ (and hence $u_2 = s'a$) for some a and $sat = u_1 s'at \neq 0$. The first term is $u_1 g_k(u_2, \dots, u_{k+1}) = \zeta_{n,a} u_1 s'at$. Like in the previous paragraph all other terms are zero.
- Similarly the last and penultimate term also cancel out.
- If an intermediate term $g_k(u_1, \dots, u_i u_{i+1}, \dots u_{k+1}) \neq 0$ then either u_i or u_{i+1} is a vertex. In the former case the previous intermediate term cancels it and in the latter case the next term does the job.

□

9.3. **Some $\mathtt{A}_\infty$-structures on $\mathtt{Gtl}\,(\mathrm{Q})$.** We will describe some specific $\mathtt{A}_\infty$-structures on $\mathtt{Gtl}\,(\mathrm{Q})$, which can be constructed inductively. Choose a map $\kappa : \mathrm{Q}_2^+ \times \mathbb{N} \to \mathbb{C}$.

For any cycle $b_1 \dots b_l$ in Q_2^+, power n, paths $p_1, \dots, p_k$ and location $i \in [1, k]$ such that

A $p_i b_1 \neq 0$ and $b_l p_{i+1} \neq 0$,
B $p_i \neq h(b_1)$ if $i = 1$ and $p_{i+1} \neq t(b_l)$ if $i + 1 = k$.

$$\mu^\kappa(p_1,\dots,p_ib_1,b_2,\dots,b_{nl-1},b_{nl}p_{i+1},\dots,p_k) := (-1)^s\kappa(b_1\dots b_l,n)\mu(p_1,\dots,p_k)$$

with sign convention $s = nl(p_1+\dots+p_i+k-i)$.



For $u > 2$ we set $\mu^\kappa(q_1,\dots,q_u) = 0$ if we cannot perform any reduction of the form above.

Lemma 9.7. *The rule above makes μ^κ_u well defined for all u.*

Proof. We have to check that if there are two positive cycles $a_r,\dots a_1$ and $b_s\dots b_1$ that can be used to reduce the product, the end result will not depend on the order of the reduction. If we look at an entry that is equal to an arrow, we can uniquely identify the cycle and the power we need to reduced this arrows away. This is because each arrow sits in just one positive cycle and property B will determine the power.

If we have two cycles that can be reduced, our big product either looks like

$$\mu_{r+s+k-4}(p_1,\dots,p_ia_1,a_2,\dots,a_{r-1},a_rp_{i+1},\dots,p_jb_1,b_2,\dots b_{s-1},b_sp_{j+1},\dots,p_k)$$

or

$$\mu_{r+s+l-4}(p_1,\dots,p_ia_1,a_2,\dots,a_r,a_1p_{i+1}b_1,b_2,\dots b_{s-1},b_sp_{j+2},\dots,p_k)$$

In both cases it is clear that, up to a sign, first reducing $a_r\dots a_1$ and then $b_s\dots b_1$ gives the same result as reducing in the opposite order. To show that the signs are the same, let us look at the first $\mu_{r+s+l-4}$. If we first reduce $a_1\dots a_r$ and then $b_1\dots b_s$, we get a total sign

$$s_{tot} = r(p_1+\dots+p_i+k-i-1+s-2)+s(p_1+\dots+p_j+k-j-1).$$

If we do it the other way round we get

$$s_{tot} = s(p_1+\dots+p_i+a_1+\dots+a_r+k-j-1)+r(p_1+\dots+p_i+k-i-1).$$

Because of the construction of the gradation in section 4, $a_1\dots a_r$ has odd degree if and only if r is odd. Therefore $s(a_1+\dots+a_r)+r(s-2) = 0 \mod 2$. The second sign calculation is similar. □

Lemma 9.8. *The $\mu_u := \mu^\kappa_u$, $u > 2$ have the following properties*

(1) *$\mu(p_1,\dots,p_u)$ is homotopic to $p_1\cdots p_u$ viewed as a path in $|Q|\setminus Q_0$.*
(2) *If $\mu(p_1,\dots,p_u) = \pm q$ is not a trivial path then either $p_1 := qp'_1$ or $p_u = p'_uq$. In the first case for any q' such that $q'p'_1 \neq 0$ we have $\mu(q'p'_1,\dots,p_u) = q'\mu(p'_1,\dots,p_u)$ in the second we have $\mu(p_1,\dots,p'_uq') = \mu(p_1,\dots,p'_u)q'$ if $p'_uq' \neq 0$.*
(3) *$\mu(p_1,\dots,p_u) = 0$ if $p_ip_{i+1} \neq 0$ for some $i < u$ or in particular when p_i is trivial.*

Proof. We prove this by induction on the number of cycles we need to reduce μ_u to an ordinary product.

1 The first statement clearly holds for μ_2 and by induction for higher multiplications because any positive cycle $a_1\cdots a_l$ is contractible in $|Q|\setminus Q_0$.
2 If there is just one cycle then $\mu(p_1a_1,\dots,a_lp_2) = p_1p_2$ but a_1 and a_l sit in a different negative cycle so p_1p_2 is zero unless one of the 2 paths has length zero. It is also clear that if p_1 has nonzero length then $\mu(qp_1a_1,\dots,a_lp_2) = qp_1$ and a similar statement holds for nontrivial p_2.

If the statement holds up to k reductions then it also holds for $k+1$ reductions because the first and last path in μ after a reduction are subpaths starting from the outer ends of the first and last paths of the unreduced μ.

3 If there is just one cycle and $\mu(p_1a_1,\dots,a_lp_2) \neq 0$ then $a_ia_{i+1} = 0$. If the statement holds up to k reductions then it also holds for $k+1$ reductions because $a_ia_{i+1} = 0$ for any positive cycle.

$\square$

Proposition 9.9. *The products μ^κ turn $\mathtt{Gtl}\,(\mathrm{Q})$ into an $\mathtt{A}_\infty$-category.*

Proof. We will show that the identity $[M_k]$

$$\sum_{s+r+t=k} (-1)^{s+rt+(2-r)(p_1+\cdots+p_s)}\mu(p_1,\dots p_s,\mu_r(p_{s+1},\dots,p_{s+r}),p_{s+r+1}\dots,p_k) = 0$$

holds for all possible collections of paths $p_1,\dots,p_k$ with $t(p_i) = h(p_{i+1})$. We will do this using induction on k.

For $k \leq 3$ the identities hold because $\mathtt{Gtl}\,(\mathrm{Q})$ is an associative algebra with zero differential. Suppose now that the identity $[M_j]$ holds for all $j < k$.

Because $k > 3$, every term in M_k will have at least one higher order multiplication in it. So if there is a nonzero term, there is at least one cycle we can reduce, so we can assume that the sequence of paths looks like

$$p_1,\dots,p_ia_1,a_2,\dots,a_{nl}p_{i+1},\dots,p_k.$$

Suppose now that we have a term that is nonzero. If the inner μ is completely contained in the cycle then it can only be nonzero if $n > 1$ and μ covers a subcycle. In that case this inner μ returns a trivial path for one of the entries of the outer μ, which will then be zero.

If the inner μ overlaps partially with the cycle then we can consider two situations:

A If the outer μ is a higher multiplication we get expressions like

$$\mu(\dots,\mu(\dots,a_j),a_{j+1},\dots) \text{ or } \mu(\dots,a_j,\mu(a_{j+1},\dots),\dots).$$

In the first case $\mu(\dots,a_j)$ must evaluate to something ending in a_j otherwise we cannot reduce a_{j+1}.

If $j \geq 2$, $\mu(\dots,a_j)$ we first reduce $\mu(\dots,a_j)$ until we get to the reduction that will remove a_{j-1}. If this reduction does not remove a_j then we end up with a higher order μ that contains a trivial path which is impossible because we assumed the term was nonzero. If this reduction does remove a_j the only way we could get something nonzero is if the situation looked like

$$\mu(qa_{j+1},a_{j+2},\dots,a_j) \to \mu(q,t(a_j)) = q.$$

But as q must end in a_j we have that $qa_{j+1} = 0$. Hence if the term is nonzero we must assume $j = 1$ and then for the same reason no reduction for the inner μ can reduce a_1.

Similarly for $\mu(\dots,a_j,\mu(a_{j+1},\dots),\dots)$ we have that $j > nl-2$ and no inner reduction can reduce a_{nl}.

B If the outer μ is the ordinary multiplication then it looks like

$$\mu(\mu(\dots,a_{nl-2},a_{nl-1}),a_{nl}p_k)$$

or a similar expression with the inner μ on the right. Now reduce the inner μ until we get to the reduction that gets rid of a_{nl-2}. It must also remove a_{nl-1} because otherwise we get a higher multiplication with a trivial path. This implies that the nonzero term looks like

$$\mu(\mu(q_1a_{nl}q_2,\dots,q_ua_1,\dots,a_{nl-2},a_{nl-1}),a_{nl}p_k)$$

(The inner reduction first gets rid of $q_2,\dots,q_u$ and after that there can only be nl terms in the inner μ because otherwise reducing $a_{nl}\dots a_{nl-1}$ will give a higher multiplication with a trivial path).

If $q_2,\dots,q_u$ is nontrivial, we pick instead of $a_1\dots a_{nl}$ a cycle in there. This cycle will then fall under case $[A]$. If $q_2,\dots,q_u$ is trivial our sequence of paths looks like

$$p_1a_{nl}, a_1, a_2, \dots, a_{nl-1}, a_np_2\ (*).$$

It is easy to check that for such cases $[M_k]$ holds.

If our sequence is not of the form $(*)$ all nonzero terms fall in 4 categories.

- The inner μ contains no part of the cycle $a_1,\dots,a_{nl}$.
- The inner μ contains the whole cycle $a_1,\dots,a_{nl}$.
- The first entry of the inner μ is $a_{nl}p_{i+1}$ and the inner μ evaluates to a right multiple of a_{nl}.
- The last entry of the inner μ is p_ia_1 and the inner μ evaluates to a left multiple of a_1.

In each of these cases this term equals

$$\pm\kappa(a_1\dots a_l,n)\mu(p_1,\dots,p_s,\mu(p_{s+1},\dots,p_{s+r}),p_{s+r+1},\dots,p_k)$$

for the appropriate s,r,t. Different terms will give different simplified terms. Putting a factor $\pm\kappa(a_1\dots a_l,n)$ in front we get that the expression $[M_{k+nl-2}]$ for

$$p_1,\dots,p_ia_1,a_2,\dots,a_{nl}p_{i+1},\dots,p_k$$

is $\pm\kappa(a_1\dots a_l,n)$ times $[M_k]$ for $p_1,\dots,p_k$.

The fact that the signs match up requires some computation. In particular we need to show that the product of the sign in $[M_{k+nl-2}]$, the sign of κ and the sign in $[M_k]$ does not depend on s,r,t and the 4 different cases (note that the first case splits in 2 depending on whether the cycle comes before or after the inner μ.)

We will only do this calculation for the first case with $i<s$ (i.e. the cycle comes before μ)

$$\begin{aligned}
&\underbrace{s+(nl-2)+rt+(2-r)(p_1+\dots+p_s+a_1+\dots+a_{nl})}_{[M_{k+nl-2}]}\\
&+\underbrace{s+rt+(2-r)(p_1+\dots+p_s)}_{M_k}+\underbrace{nl(p_1+\dots+p_i)+s+t+1-i}_{\kappa}\\
&=(nl-\cancel{2})+(\cancel{2}-r)(\underbrace{a_1+\dots+a_{nl}}_{=nl})+nl(s+t+1-i+p_1+\dots+p_i)\\
&=nl(\underbrace{r+s+t}_{=k}+1-i+p_1+\dots+p_i)\quad\mod 2
\end{aligned}$$

and the second case (when the cycle is inside the inner μ: $s+t>i>s$)

$$\begin{aligned}
&\underbrace{s+(r+nl-\cancel{2})t+(\cancel{2}-r-nl+\cancel{2})(p_1+\dots+p_s)}_{[M_{k+nl-2}]}\\
&+\underbrace{s+rt+(\cancel{2}-r)(p_1+\dots+p_s)}_{M_k}+\underbrace{nl(p_{s+1}+\dots+p_i)+r-i+s}_{\kappa}\\
&=nlt-nl(p_1+\dots+p_s)+nl(r-i+s+p_{s+1}+\dots+p_i)\\
&=nl(\underbrace{r+s+t}_{=k}+1-i+p_1+\dots+p_i)\quad\mod 2
\end{aligned}$$

□

Given a map $\rho:\mathrm{RQ}_1\to\mathbb{C}^*$ we get an autoequivalence of the category $\mathtt{Fuk}\mathrm{Q}$

$$f_\rho:\mathtt{Gtl}\,(\mathrm{Q})\to\mathtt{Gtl}\,(\mathrm{Q}):a\mapsto\rho(a)a.$$

This autoequivalence turns an $\mathtt{A}_\infty$-structure defined by κ into a new one, which is defined by $\kappa':\mathrm{Q}_2^+\to\mathbb{C}:a_1\dots a_k\mapsto\kappa(a_1\dots a_k)\rho(a_1)\cdots\rho(a_k)$.

Lemma 9.10. *Let* $\kappa_1, \kappa_2 : \mathrm{Q}_2^+ \times \mathbb{N} \to \mathbb{C}$ *be two maps for which* $\kappa_i(c,n) = 0$ *as soon as* $n > 1$*. The two* $\mathtt{A}_\infty$*-structures defined by* κ_1 *and* κ_2 *are isomorphic if* $\forall c \in \mathrm{R}Q_2^+ :$ $\kappa_1(c) = 0 \iff \kappa_2(c) = 0$*.*

Proof. Choose an arrow a_c in each positive cycle c with $\kappa_1(c,1) \neq 0$ and define $\rho(a_c) = \kappa_2(c,1)\kappa_1(c,1)^{-1}$ and set all other $\rho(a) = 1$. This gives an isomorphism between the two $\mathtt{A}_\infty$-structures. □

9.4. **Identifying $\mathtt{A}_\infty$-structures on $\mathtt{Gtl}\,(\mathrm{Q})$.**

Lemma 9.11. *If* μ *is an* $\mathtt{A}_\infty$*-structure on* $\mathtt{Gtl}\,(\mathrm{Q})$ *and* $a_1, \dots a_l$ *is a positive cycle and we can find a* $\kappa \in \mathbb{C}$ *such that*

$$\mu(a_1, \dots, a_{nl}) = \kappa h(a_1)$$

then

$$\mu(a_i, \dots, a_{nl+i-1}) = \kappa h(a_i).$$

Proof. This follows straight from the identity $[M_{nl+1}]$ applied to the paths $a_1, \dots, a_{nl}, a_1$:

$$\mu(a_1, \mu(a_2, \dots, a_{nl})) - \mu(\mu(a_1, \dots, a_{nl}), a_1) = 0.$$

□

Theorem 9.12. *Let* $\mathtt{Gtl}\,(\mathrm{Q})$ *be the gentle category coming from a rectified dimer* Q*. Let* μ *be an* $\mathtt{A}_\infty$*-structure on* $\mathtt{Gtl}\,(\mathrm{Q})$ *such that for every positive cycle* $a_1 \dots a_l$ *we have*

$$\mu(a_i, \dots, a_j) = \begin{cases} \kappa(a_1 \dots a_l, n) h(a_i) & j - i + 1 = nl \\ 0 & \textit{otherwise,} \end{cases}$$

where

$$\kappa : \mathrm{Q}_2^+ \times \mathbb{N} \to \mathbb{C} : (a_1 \dots a_l, n) \mapsto \mu(a_1, \dots, a_{nl}).$$

is a map. Then μ *is* $\mathtt{A}_\infty$*-isomorphic to* μ^κ*.*

Proof. In order to prove that μ^κ and μ are isomorphic we construct an $\mathtt{A}_\infty$-functor $\mathcal{F}$ with $\mathcal{F}_1 = \mathbf{1}$. We show that we can do this by constructing the $\mathcal{F}_i$ one at a time. Suppose we have constructed $\mathcal{F}_i$ for $i < r$.

Now look at the identity $[F_n]$. We already know it is of the form

$$d\mathcal{F}_n = \Psi$$

with $d\Psi = 0$. In order to show that we can find an $\mathcal{F}_n$ we have to show that $\Psi \in \mathrm{Im} d$ or equivalently that the homology class of Ψ is zero. If we prove that $\Psi(a_i, \dots, a_{i+nl-1})$ and $\Psi(a_i, \dots, a_{i+nl})$ contain no length 0 or length 1 terms, then by lemma 9.5 we are done.

The terms in the expression $\Psi(a_i, \dots, a_r)$ are of the forms

- $\mu(a_i, \dots, a_{r-i}) - \mu^\kappa(a_i, \dots, a_{r-i})$. This expression contains no length 0 or length 1 terms by lemma 9.11.
- $\mathcal{F}(a_i, \dots, a_j, \mu(a_{j+1}, \dots, a_u), a_{u+1}, \dots, a_r)$. Such a term is zero because by the condition in the theorem $\mu(a_{j+1}, \dots, a_u)$ can only be zero or a multiple of $h(a_{j+1})$. But then $\mathcal{F}$ contains an idempotent so it is zero by lemma 9.13.
- $\mu^\kappa(\mathcal{F}(a_1, \dots, a_{i_1}), \dots, \mathcal{F}(a_{i_j}, \dots, a_r))$. If $\mathcal{F}(a_i, \dots, a_{i+s})$ is nonzero then by lemma 4.3, s must be $-2, -1, 0 \mod l$.
  - If it is $-2 \mod l$ then $\mathcal{F}(a_i, \dots, a_{i+s})$ sits in $\mathtt{Hom}(h(a_i), t(a_1))$, so all its terms have higher length.
  - If it is $-1 \mod l$ then the degree of $\mathcal{F}$ (s) and the degree of $a_i, \dots, a_{i+s}$ ($s+1$) imply that $\mathcal{F}(a_i, \dots, a_{i+s})$ has odd degree, which contradicts lemma 4.3.
  - If it is $0 \mod l$ then $\mathcal{F}(a_i, \dots, a_{i+s})$ sits in $\mathtt{Hom}(t(a_i), h(a_i))$ and can have a term which is a multiple of a_i but we can always add an element in $\ker d$ to f so by lemma 9.6 we can assume this term is zero.

  In other words all terms are zero or have higher length. By construction μ^κ of such an expression is zero, because there is no position where we can start to reduce it.

□

Lemma 9.13. *Let* $(\mathcal{F}_i)$ *be a partial* $\mathtt{A}_\infty$*-functor between 2* $\mathtt{A}_\infty$*-structures* μ^1 *and* μ^2*, for which* $\mathcal{F}_1$ *is the identity functor. For each object* X *we have that*

$$\mathcal{F}_k(u_1, \dots u_{l-1}, \mathbf{1}_X, u_{l+1}, \dots, u_k) = 0$$

where the u_i *are any maps.*

Proof. We first show that $\mathcal{F}_k(1_X, \dots, 1_X) = 0$. If k is even this is true because $\deg \mathcal{F} = 1-k$ is odd and there are no odd elements in $\mathtt{Hom}(X,X)$. If k is odd then we use induction: suppose it holds for $i < k$. Now

$$\begin{aligned} d\mathcal{F}_k(1_X, \dots, 1_X) &= \underbrace{1_X\mathcal{F}_k(1_X, \dots, 1_X) - \mathcal{F}_k(1_X \cdot 1_X, \dots, 1_X) \cdots \pm \mathcal{F}_k(1_X, \dots, 1_X)1_X}_{k+2 \text{ terms}} \\ &= \mathcal{F}_k(1_X, \dots, 1_X) \end{aligned}$$

because the number of terms is odd. But on the other hand we know that because $\mathcal{F}$ is a partial functor the following identity holds:

$$\begin{aligned} d\mathcal{F}_k(1_X, \dots, 1_X) = &\sum_{r+s+t=k} \pm\mathcal{F}(\underbrace{1_X, \dots, 1_X}_{\times r}, \mu^1(\underbrace{1_X, \dots, 1_X}_{\times s}), \underbrace{1_X, \dots, 1_X}_{\times t}) \\ &- \sum_{i_1+\dots+i_u=k} \pm\mu^2(\mathcal{F}(\underbrace{1_X, \dots, 1_X}_{\times i_1}), \dots, \mathcal{F}(\underbrace{1_X, \dots, 1_X}_{\times i_u})). \end{aligned}$$

each of these terms is zero because both the $\mu^i(1_X, \dots, 1_X)$ are zero and $\mathcal{F}(1_X, \dots, 1_X) = 0$ by induction.

To prove that $\mathcal{F}_k(u_1, \dots u_{l-1}, 1_X, u_{l+1}, \dots, u_k) = 0$ we use double induction: on k and on the number of entries u_i that are not the identity. Now

$$\begin{aligned} d\mathcal{F}_k(u_1, \dots u_{l-1}, 1_X, 1_X, u_{l+1}, \dots, u_k) = &+ u_1\mathcal{F}_k(u_2, \dots, 1_X, 1_X, u_k) - \dots \\ &\pm \mathcal{F}_k(u_1, \dots u_{l-1} \cdot 1_X, 1_X, u_{l+1}, \dots, u_k) \\ &\mp \mathcal{F}_k(u_1, \dots u_{l-1}, 1_X \cdot 1_X, u_{l+1}, \dots, u_k) \\ &\pm \mathcal{F}_k(u_1, \dots u_{l-1}, 1_X, 1_X \cdot u_{l+1}, \dots, u_k) \\ &+ \dots \mathcal{F}_k(u_2, \dots, 1_X, 1_X, \dots, u_{k-1})u_k \end{aligned}$$

The first and last line are zero by induction because they have even more entries that are the identity, the intermediate lines contain 3 terms with alternating sign that are equal to

$$\mathcal{F}_k(u_1, \dots u_{l-1}, 1_X, u_{l+1}, \dots, u_k).$$

Again we can look at the identity for the partial $\mathtt{A}_\infty$-functor

$$d\mathcal{F}_k(\dots) = \sum_{r+s+t=k} \pm\mathcal{F}(\dots, \mu^1(\dots), \dots) - \sum_{i_1+\dots+i_u=k} \pm\mu^2(\mathcal{F}(\dots), \dots, \mathcal{F}(\dots)).$$

Each term of it is zero: either because it contains an f_i with $i < k$ with a 1_X entry or a μ^1 or μ^2 with a 1_X entry. □

REFERENCES

1. M. Abouzaid, *Morse homology, tropical geometry, and homological mirror symmetry for toric varieties*, Selecta Math. 15 (2009), 189270.
2. M. Abouzaid, P. Seidel, *An open string analogue of Viterbo functoriality*, Geom. Topol. 14 (2010), 627718.
3. I. Assem, T. Brüstle, G. Charbonneau-Jodoin, and P. Plamondon, *Gentle algebras arising from surface triangulations*, Algebra Number Theory 4 (2010), no. 2, 201229.
4. I. Assem and A.Skowrónski, Iterated tilted algebras of type A n, Math. Z. 195 (1987), no. 2, 269-290.
5. D. Auroux, L. Katzarkov, D. Orlov, *Mirror symmetry for weighted projective planes and their noncommutative deformations*, Ann. Math. 167 (2008), 867943.
6. D. Auroux, L. Katzarkov, D. Orlov, *Mirror symmetry for Del Pezzo surfaces: Vanishing cycles and coherent sheaves*, Inventiones Math. 166 (2006), 537582.
7. M. Abouzaid, D. Auroux, A. Efimov, L. Katzarkov, D. Orlov *Homological mirror symmetry for punctured spheres*, `arXiv:1103.4322`

8. M. Bardzell, *The Alternating Syzygy Behavior of Monomial Algebras*, Journal of Algebra 188, 69-89 (1997).
9. R. Bocklandt, *Calabi-Yau Algebras and Quiver Polyhedra*, `arXiv:0905.0232`
10. R. Bocklandt, *Consistency conditions for dimer models*, `arXiv:1104.1592`
11. N. Broomhead, *Dimer models and Calabi-Yau algebras*, Memoirs of the AMS, 2011, 86 pp. MEMO/215/1011 `arXiv:0901.4662`
12. B. Davison, *Consistency conditions for brane tilings*, `arXiv:0812.4185`
13. Bo. Feng, Y. He, K. Kennaway, C. Vafa *Dimer Models from Mirror Symmetry and Quivering Amoebae* Adv. Theor. Math. Phys. Volume 12, Number 3 (2008), 489-545.
14. S. Franco, A. Hanany, K. D. Kennaway, D. Vegh, B. Wecht, *Brane Dimers and Quiver Gauge Theories*, JHEP 0601 (2006) 096, `hep-th/0504110`.
15. V. Ginzburg, *Calabi-Yau algebras*, `math/0612139`.
16. A. Hanany, C. P. Herzog, D. Vegh, *Brane Tilings and Exceptional Collections*, J. High Energy Phys. 2006, no. 7, 001, 44 pp.
17. A. Hanany, K. D. Kennaway, *Dimer models and toric diagrams*, `hep-th/0602041`.
18. A. Hanany, D. Vegh, *Quivers, tilings, branes and rhombi*, J. High Energy Phys. 2007, no. 10, 029, 35 pp.
19. A. Hatcher, *On triangulations of surfaces*, Topology and its Applications 40 (1991), 189-194.
20. K. Hori, C. Vafa, *Mirror symmetry*, `arXiv:hep-th/0002222`.
21. D. Gulotta, *Properly ordered dimers, R-charges, and an efficient inverse algorithm*, JHEP0810:014, 2008 `arXiv:0807.3012`
22. A. Ishii, K. Ueda, *On moduli spaces of quiver representations associated with brane tilings* Higher dimensional algebraic varieties and vector bundles, 127141, RIMS Kokyuroku Bessatsu, B9, Res. Inst. Math. Sci. (RIMS), Kyoto, 2008
23. A. Ishii, K. Ueda, *Dimer models and exceptional collections*, `arXiv:0911.4529`
24. A. Ishii, K. Ueda, *A note on consistency conditions on dimer models* `arXiv:1012.5449`.
25. T. Kadeishvili *The algebraic structure in the homology of an A_∞-algebra.* (Russian) Soobshch. Akad. Nauk Gruzin. SSR 108 (1982), no. 2, 249252 (1983)
26. L. Katzarkov. *Birational geometry and homological mirror symmetry*. In Real and complex singularities, pages 176206. World Sci. Publ., 2007.
27. B. Keller, *Introduction to A-infinity algebras and modules*. Homology Homotopy Appl. 3 (2001), 135.
28. K. D. Kennaway, *Brane tilings*, Internat. J. Modern Phys. A 22 (2007), no. 18, 29773038. `hep-th/0710.1660`.
29. R. Kenyon, *An introduction to the dimer model*, School and Conference on Probability Theory, 267304, ICTP Lect. Notes, XVII, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004
30. M. Kontsevich, *Homological algebra of mirror symmetry*, Proc. International Congress of Mathematicians (Zürich, 1994), Birkhäuser, Basel, 1995, pp. 120139.
31. M. Kontsevich; Y. Soibelman *Deformations of algebras over operads and the Deligne conjecture*, Conférence Moshe Flato 1999, Vol. I (Dijon), Math. Phys. Stud., vol. 21, Kluwer Acad. Publ., Dordrecht, 2000, pp. 255307.
32. M. Kontsevich, Y. Soibelman *Notes on A_∞-algebras, A_∞-categories and non-commutative geometry.* Homological mirror symmetry, 153219, Lecture Notes in Phys., 757, Springer, Berlin, 2009.
33. S. Mozgovoy, *Crepant resolutions and brane tilings I: Toric realization* `arXiv:0908.3475`
34. M. Bender, S. Mozgovoy, *Crepant resolutions and brane tilings II: Tilting bundles* `arXiv:0909.2013`
35. S. Mozgovoy, M. Reineke, *On the noncommutative Donaldson-Thomas invariants arising from brane tilings*, Adv. Math. 223 (2010), no. 5, 15211544
36. D. Orlov. *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, 2003.
37. D. Orlov. *Triangulated categories of singularities and equivalences between Landau-Ginzburg models*, 2005.
38. A. Quintero Vélez, *McKay correspondence for Landau-Ginzburg models.* Commun. Number Theory Phys. 3 (2009), no. 1, 173208.
39. P. Seidel, Fukaya categories and Picard-Lefschetz theory, Zurich Lect. in Adv. Math., European Math. Soc., Zürich, 2008.
40. M. Van den Bergh, *Non-commutative crepant resolutions*, The Legacy of Niels Hendrik Abel, Springer, pp. 749-770, 2002.

RAF BOCKLANDT, SCHOOL OF MATHEMATICS AND STATISTICS, HERSCHEL BUILDING, NEWCASTLE UNIVERSITY, NEWCASTLE UPON TYNE, NE1 7RU, UK
E-mail address: `raf.bocklandt@gmail.com`